



5-13-2014

Cytochrome P450S and Uses Thereof

Joe Chappell

University of Kentucky, chappell@uky.edu

Lyle F. Ralston

Right click to open a feedback form in a new tab to let us know how this document benefits you.

Follow this and additional works at: https://uknowledge.uky.edu/pss_patents



Part of the [Plant Sciences Commons](#)

Recommended Citation

Chappell, Joe and Ralston, Lyle F., "Cytochrome P450S and Uses Thereof" (2014). *Plant and Soil Sciences Faculty Patents*. 16.
https://uknowledge.uky.edu/pss_patents/16

This Patent is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in Plant and Soil Sciences Faculty Patents by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.



US008722363B2

(12) **United States Patent**
Chappell et al.(10) **Patent No.:** **US 8,722,363 B2**(45) **Date of Patent:** ***May 13, 2014**(54) **CYTOCHROME P450S AND USES THEREOF**(71) Applicant: **Lyle F. Ralston**, São Paulo (BR)(72) Inventors: **Joseph Chappell**, Lexington, KY (US);
Lyle F. Ralston, São Paulo (BR)(73) Assignee: **University of Kentucky Research**
Foundation, Lexington, KY (US)(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.This patent is subject to a terminal dis-
claimer.(21) Appl. No.: **13/986,446**(22) Filed: **May 3, 2013**(65) **Prior Publication Data**

US 2013/0330793 A1 Dec. 12, 2013

Related U.S. Application Data(63) Continuation of application No. 13/199,349, filed on
Aug. 26, 2011, now Pat. No. 8,445,231, which is a
continuation of application No. 12/182,000, filed on
Jul. 29, 2008, now Pat. No. 8,263,362, which is a
continuation of application No. 10/097,559, filed on
Mar. 8, 2002, now Pat. No. 7,405,057.(60) Provisional application No. 60/274,421, filed on Mar.
9, 2001, provisional application No. 60/275,597, filed
on Mar. 13, 2001.(51) **Int. Cl.****C12N 1/00** (2006.01)
C12N 1/06 (2006.01)
C12N 5/00 (2006.01)
C12N 5/07 (2010.01)(52) **U.S. Cl.**USPC **435/69.1**; 435/468; 435/348; 435/252.1;
435/252.2; 435/155; 435/419(58) **Field of Classification Search**

None

See application file for complete search history.

(56) **References Cited****U.S. PATENT DOCUMENTS**5,589,619 A 12/1996 Chappell et al. 800/205
5,672,487 A 9/1997 Schweden et al. 435/69.1
5,741,674 A 4/1998 Schweden et al. 435/69.1
5,766,911 A 6/1998 Koike et al. 435/193
5,824,774 A 10/1998 Chappell et al. 530/350
5,981,843 A 11/1999 Chappell et al. 800/301
5,994,114 A 11/1999 Croteau et al. 435/232
6,072,045 A 6/2000 Chappell et al. 536/23.1
6,100,451 A 8/2000 Chappell et al. 800/298
6,117,649 A 9/2000 Bellamine et al. 435/25
6,194,185 B1 2/2001 Croteau et al. 435/189
6,331,660 B1 12/2001 Chomet et al. 800/278
6,368,837 B1 4/2002 Gatenby et al. 435/146
6,468,772 B1 10/2002 Chappell et al. 435/183
6,495,354 B2 12/2002 Chappell et al. 435/1836,531,303 B1 3/2003 Millis et al. 435/155
6,559,297 B2 5/2003 Chappell et al. 536/23.1
6,569,656 B2 5/2003 Chappell et al. 435/183
6,645,762 B2 11/2003 Chappell et al. 435/325
6,689,593 B2 2/2004 Millis et al. 435/155
6,890,752 B2 5/2005 Chappell et al. 435/325
7,186,891 B1 3/2007 Chappell et al. 800/298
7,405,057 B2 * 7/2008 Chappell et al. 435/69.1
7,442,785 B2 10/2008 Chappell et al. 536/23.6
7,622,614 B2 11/2009 Julien et al. 568/327
8,106,260 B2 1/2012 Chappell et al. 800/298
8,124,811 B2 2/2012 Julien et al. 568/367
8,192,950 B2 6/2012 Chappell et al. 435/41
8,263,362 B2 * 9/2012 Chappell et al. 435/69.1
8,354,504 B2 1/2013 Chappell et al. 530/379
8,362,309 B2 1/2013 Julien et al. 583/360
8,445,231 B2 * 5/2013 Chappell et al. 435/69.1
8,481,286 B2 7/2013 Julien et al. 435/69.1
2003/0166255 A1 9/2003 Chappell 435/252.3
2004/0078840 A1 4/2004 Chappell et al. 800/278
2006/0218661 A1 9/2006 Chappell et al. 800/278
2007/0231861 A1 10/2007 Millis et al. 435/69.1
2007/0238157 A1 10/2007 Millis et al. 435/166
2007/0238159 A1 10/2007 Millis et al. 435/252.33
2007/0238160 A1 10/2007 Millis et al. 435/252.33
2007/0254354 A1 11/2007 Millis et al. 435/252.33

(Continued)

FOREIGN PATENT DOCUMENTSJP 2000-511404 9/2000
WO 96/36697 11/1996

(Continued)

OTHER PUBLICATIONSInternational Search Report, issued Dec. 17, 2002, in connection with
corresponding International Patent Application No. PCT/US02/
06912, 4 pages.International Preliminary Examination Report, issued Apr. 24, 2003,
in connection with corresponding International Patent Application
No. PCT/US02/06912, 4 pages.Partial European Search Report, issued Jun. 16, 2004, in connection
with corresponding European Patent Application No. 02709797.1, 7
pages.Office Action, issued Jul. 1, 2004, in connection with corresponding
U.S. Appl. No. 10/097,559, 10 pages.Supplemental European Search Report, issued Sep. 8, 2004, in con-
nection with corresponding European Patent Application No.
02709797.1, 6 pages.

(Continued)

Primary Examiner — Medina A Ibrahim(74) *Attorney, Agent, or Firm* — McKenna Long & Aldridge
LLP; Stephanie Seidman(57) **ABSTRACT**The invention features isolated cytochrome P450 polypep-
tides and nucleic acid molecules, as well as expression vec-
tors and transgenic plants containing these molecules. In
addition, the invention features uses of such molecules in
methods of increasing the level of resistance against a disease
caused by a plant pathogen in a transgenic plant, in methods
for producing altered compounds, for example, hydroxylated
compounds, and in methods of producing isoprenoid com-
pounds.**29 Claims, 11 Drawing Sheets**

(56)

References Cited

U.S. PATENT DOCUMENTS

2008/0178354	A1	7/2008	Chappell et al.	800/298
2008/0233622	A1	9/2008	Julien et al.	435/148
2010/0035329	A1	2/2010	Millis et al.	435/254.2
2010/0120110	A1	5/2010	Chappell	435/166
2010/0129306	A1	5/2010	Julien et al.	424/65
2010/0151519	A1	6/2010	Julien et al.	435/69.1
2010/0151555	A1	6/2010	Julien et al.	435/193
2010/0216186	A1	8/2010	Chappell et al.	435/69.1
2011/0081703	A1	4/2011	Chappell et al.	435/193
2011/0318797	A1	12/2011	Chappell et al.	435/155
2012/0129235	A1	5/2012	Julien et al.	435/166
2012/0196340	A1	8/2012	Chappell et al.	435/148
2013/0071877	A1	3/2013	Chappell et al.	435/41
2013/0122560	A1	5/2013	Julien et al.	435/148
2013/0236943	A1	9/2013	Julien et al.	435/166

FOREIGN PATENT DOCUMENTS

WO	WO 97/37664	10/1997
WO	WO 97/38571	10/1997
WO	WO 97/38703	10/1997
WO	WO 00/17327	3/2000
WO	WO 02/072758	9/2002
WO	WO 2010/019696	2/2010

OTHER PUBLICATIONS

Examination Report, issued Sep. 19, 2005, in connection with corresponding European Patent Application No. 02709797.1, 6 pages.

Response to Examination Report, filed Feb. 9, 2006, in connection with corresponding European Patent Application No. 02709797, 11 pages.

Office Action, issued Mar. 13, 2006, in connection with corresponding U.S. Appl. No. 10/097,559, 9 pages.

Office Action, issued Oct. 12, 2006, in connection with corresponding U.S. Appl. No. 10/097,559, 12 pages.

Office Action, issued May 1, 2007, in connection with corresponding U.S. Appl. No. 10/097,559, 10 pages.

Examination Report, issued May 28, 2007, in connection with corresponding Canadian Patent Application No. 2,440,278, 4 pages.

Response to Examination Report, filed Nov. 28, 2007, in connection with corresponding Canadian Patent Application No. 2,440,278, 13 pages.

Official Action, issued Dec. 18, 2007, in connection with corresponding Japanese Patent Application No. 2008-28198, 15 pages.

Examination Report, issued Apr. 11, 2008, in connection with corresponding European Patent Application No. 02709797.1, 6 pages.

Examination Report, issued May 7, 2008, in connection with corresponding Canadian Patent Application No. 2,440,278, 3 pages.

Response to Examination Report, filed Nov. 6, 2008, in connection with corresponding Canadian Patent Application No. 2,440,278, 11 pages.

Response to Examination Report, filed Dec. 18, 2008, in connection with corresponding European Patent Application No. 02709797.1, 11 pages.

Office Action, issued Apr. 14, 2010, in connection with corresponding U.S. Appl. No. 12/182,000, 16 pages.

Examination Report, issued Aug. 2, 2010, in connection with corresponding Canadian Patent Application No. 2,440,278, 3 pages.

Intent to Grant, issued Dec. 29, 2010, in connection with corresponding European Patent Application No. 02709797.1, 5 pages.

Office Action, issued Jan. 6, 2011, in connection with corresponding U.S. Appl. No. 12/182,000, 10 pages.

Decision to Grant, issued May 19, 2011, in connection with corresponding European Patent Application No. 02709797.1, 1 page.

Appeal Decision, issued Oct. 21, 2011, in connection with corresponding Japanese Patent Application No. 2002-571814, 17 pages.

Response to Examination Report, submitted Feb. 2, 2012, in connection with corresponding Canadian Patent Application No. 2,440,278, 36 pages.

Office Action, issued Apr. 13, 2012, in connection with corresponding U.S. Appl. No. 13/199,349, 18 pages.

Examination Report, issued Nov. 20, 2012, and received Nov. 23, 2012, in connection with corresponding Canadian Patent Application No. 2,440,278, 7 pages.

Akiyoshi-Shibata et al., "Further oxidation of hydroxycalcidiol by calcidiol 24-hydroxylase. A study with the mature enzyme expressed in *Escherichia coli*," Eur. J. Biochem. 224:335-343 (1994).

Allylix, "Protein engineering and chemobiosynthesis to produce novel sesquiterpenoids," Presentation at BIO World Congress on Industrial Biotechnology & Bioprocessing, Washington, D.C., Jun. 28, 2010, 19 pages.

An et al., "Functional analysis of the 3' control region of the potato wound-inducible proteinase inhibitor II gene," Plant Cell 1:115-122 (1989).

An et al., "Organ-specific and developmental regulation of the nopaline synthase promoter in transgenic tobacco plants," Plant Physiol. 88:547-552 (1988).

Andersson et al., "Physiology and molecular genetics of 17 beta-hydroxysteroid dehydrogenases," Steroids 62(1):143-147 (1997).

Back et al., "Cloning and bacterial expression of sesquiterpene cyclase, a key branch point enzyme for the synthesis of sesquiterpenoid phytoalexin capsidiol in UV-challenged leaves of *Capsicum annuum*," Plant Cell. Physiol. 39:899-904 (1998).

Back et al., "Expression of a plant sesquiterpene cyclase gene in *Escherichia coli*," Arch. Biochem. Biophys. 315:527-532 (1994).

Back et al., "Cloning and bacterial expression of a sesquiterpene cyclase from *Hyoscyamus muticus* and its molecular comparison to related terpene cyclases," J. Biol. Chem. 270:7375-7381 (1995).

Back et al., "Identifying functional domains within terpene cyclases using a domain-swapping strategy," Proc. Natl. Acad. Sci. U.S.A. 93:6841-6845 (1996).

Beckman et al., "Human 25-hydroxyvitamin D3-24-hydroxylase, a multicatalytic enzyme," Biochem. 35:8465-8472 (1996).

Boddupalli et al., "Fatty acid monooxygenation by P450BM-3: product identification and proposed mechanisms for the sequential hydroxylation reactions," Arch. Biochem. Biophys. 292:20-28 (1992).

Bozak et al., "Sequence analysis of ripening-related cytochrome P-450 cDNAs from avocado fruit," Proc. Natl. Acad. Sci. U.S.A. 87:3904-3908 (1990).

Bustos et al., "Regulation of β -glucuronidase expression in transgenic tobacco plants by an A/T-rich, cis-acting sequence found upstream of a French bean β -phaseolin gene," Plant Cell 1:839-853 (1989).

Callis et al., "Heat inducible expression of a chimeric maize hsp70CAT gene in maize protoplasts," Plant Physiol. 88:965-968 (1988).

Callis et al., "Introns increase gene expression in cultured maize cells," Genes Dev. 1:1183-1200 (1987).

Cane et al., "Aristolochene biosynthesis and enzymatic cyclization of farnesyl pyrophosphate," J. Am. Chem. Soc. 111:8914-8916 (1989).

Cane et al., "Trichodiene biosynthesis and the stereochemistry of the enzymatic cyclization of farnesyl pyrophosphate," Bioorg. Chem. 13:246-265 (1985).

Cane, "Enzymatic formation of sesquiterpenes," Chem. Rev. 90:1089-1103 (1990).

Chappell et al., "Accumulation of capsidiol in tobacco cell cultures treated with fungal elicitor," Phytochem. 26:2259-2260 (1987).

Chappell et al., "Elicitor-inducible 3-hydroxy-3-methylglutaryl coenzyme A reductase activity is required for sesquiterpene accumulation in tobacco cell suspension cultures," Plant Physiol. 97:693-698 (1991).

Chappell et al., "Is the reaction catalyzed by 3-hydroxy-3-methylglutaryl coenzyme A reductase a rate-limiting step for isoprenoid biosynthesis in plants," Plant Physiol. 109:1337-1343 (1995).

Chappell, "Biochemistry and molecular biology of the isoprenoid biosynthetic pathway in plants," Annu. Rev. Plant Physiol. Plant Mol. Biol. 46:521-547 (1995).

Chappell et al., "Induction of sesquiterpenoid biosynthesis in tobacco cell suspension cultures by fungal elicitor," Plant Physiol. 85:469-473 (1987).

Chappell, "The biochemistry and molecular biology of isoprenoid metabolism," Plant Physiol. 107:1-6 (1995).

(56)

References Cited

OTHER PUBLICATIONS

- Chappell, "The genetics and molecular genetics of terpene and sterol origami," *Curr. Opin. Plant Biol.* 5:151-157 (2002).
- Chapple, "Molecular-genetic analysis of plant cytochrome P450-dependent monooxygenases," *Ann. Rev. Plant Physiol. Plant Mol. Biol.* 49:311-353 (1998).
- Chen et al., "Cloning, expression and characterization of (+)- δ -cadinene synthase: a catalyst for cotton phytoalexin biosynthesis," *Arch. Biochem. Biophys.* 324:255-266 (1995).
- Chiu et al., "Engineered GFP as a vital reporter in plants," *Curr. Biol.* 6:325-330 (1996).
- Clark et al., "Spatially distinct expression of two new cytochrome P450s in leaves of *Nepeta racemosa*: identification of a trichome-specific isoform," *Plant Mol. Biol.* 33:875-885 (1997).
- Coolbaugh et al., "Studies on the specificity and site of action of a α -cyclopropyl- α -[p-methoxyphenyl]-5-pyrimidine methyl alcohol (amcymidol), a plant growth regulator," *Plant Physiol.* 62:571-576 (1978).
- Cooper et al., "Mutagenicity of nitrosamines in methyltransferase-deficient strains of *Salmonella typhimurium* coexpressing human cytochrome P450 2E1 and reductase," *Mutat. Res.* 454:45-52 (2000).
- Deguerry et al., "The diverse sesquiterpene profile of patchouli, *Pogostemon cablin*, is correlated with a limited number of sesquiterpene synthases," *Arch. Biochem. Biophys.* 454:123-136 (2006).
- Dekeyser et al., "Transient gene expression in intact and organized rice tissues," *Plant Cell* 2:591-602 (1990).
- Devarenne et al., "Molecular characterization of tobacco squalene synthase and regulation in response to fungal elicitor," *Arch. Biochem. Biophys.* 349:205-215 (1998).
- Devarenne et al., "Regulation of squalene synthase, a key enzyme of sterol biosynthesis, in tobacco," *Plant Physiol.* 129:1095-1106 (2002).
- Diener et al., "Sterol methyltransferase 1 controls the level of cholesterol in plants," *Plant Cell* 12:853-870 (2000).
- Dietz et al., "Nucleotide sequences of subunit E of the vacuolar proton-ATPase of *Spinacia oleracea* (Accession No. X96785) and *Arabidopsis thaliana* (Accession No. X921117)," (*Plant Gene Register* PGR 96-037) *Plant Physiol.* 111:652 (1996).
- Dong et al., "Coexpression of mammalian cytochrome P450 and reductase in *Escherichia coli*," *Arch. Biochem. Biophys.* 327:254-259 (1996).
- Draper et al., "Ti plasmid homologous sequences present in tissues from *Agrobacterium* plasmid-transformed *Petunia* protoplasts," *Plant Cell Physiol.* 23:451-458 (1982).
- Facchini et al., "Gene family for an elicitor-induced sesquiterpene cyclase in tobacco," *Proc. Natl. Acad. Sci. U.S.A.* 89:11088-11092 (1992).
- Fahrendorf et al., "Stress responses in alfalfa (*Medicago sativa* L.) XVIII: Molecular cloning and expression of the elicitor-inducible cinnamic acid 4-hydroxylase cytochrome P450," *Arch. Biochem. Biophys.* 305:509-515 (1993).
- Fang et al., "Multiple cis regulatory elements for maximal expression of the cauliflower mosaic virus 35S promoter in transgenic plants," *Plant Cell* 1:141-150 (1989).
- Freeman et al., "A comparison of methods for plasmid delivery into plant protoplasts," *Plant Cell Physiol.* 25:1353-1365 (1984).
- Frohman et al., "Rapid production of full-length cDNAs from rare transcripts: amplification using a single gene-specific oligonucleotide primer," *Proc. Natl. Acad. Sci. U.S.A.* 85:8998-9002 (1988).
- Fromm et al., "An octopine synthase enhancer element directs tissue-specific expression and binds ASF-1, a factor from tobacco nuclear extracts," *Plant Cell* 1:977-984 (1989).
- Fromm et al., "Stable transformation of maize after gene transfer by electroporation," *Nature* 319:791-793 (1986).
- Gasser et al., "Genetically engineering plants for crop improvement," *Science* 244:1293-1299 (1989).
- Genbank Accession No. A35867 [online], "Cytochrome P450," Published on Sep. 26, 2000 [retrieved on Mar. 27, 2002] [retrieved from the Internet<URL:ncbi.nlm.nih.gov/nuccore/AB015762>] [1 page].
- Genbank Accession No. AAC39505 [online], "GA3 [*Arabidopsis thaliana*]," Published on Jul. 26, 1998 [retrieved on Mar. 27, 2002] [retrieved from the Internet:<URL:ncbi.nlm.nih.gov/protein/AAC39505>] [1 page].
- Genbank Accession No. AAD44150 [online], "cytochrome p450 [*Mentha spicata*]," Published on May 1, 2001 [retrieved on Mar. 27, 2002] [retrieved from the Internet<URL:ncbi.nlm.nih.gov/protein/AAD44150>] [1 page].
- Genbank Accession No. AAD44151 [online], "cytochrome p450 isoform PM17 [*Mentha x piperita*]," Published on May 1, 2001 [retrieved on Mar. 27, 2002] [retrieved from the Internet<URL:ncbi.nlm.nih.gov/protein/AAD44151>] [1 page].
- Genbank Accession No. AB015762 [online], "*Nicotiana tabacum* CYP82E1 mRNA for cytochrome P450, complete cds," Published on Sep. 26, 2000 [retrieved on Mar. 27, 2002] [retrieved from the Internet:<URL:ncbi.nlm.nih.gov/nuccore/AB015762>] [1 page].
- Genbank Accession No. CAA70575 [online], "cytochrome P450 [*Nepeta racemosa*]," Published on Sep. 9, 2004 [retrieved on Mar. 27, 2002] [retrieved from the Internet:<URL:ncbi.nlm.nih.gov/protein/CAA70575>] [1 page].
- Genbank Accession No. CAC24711 [online], "cytochrome P450 [*Solanum tuberosum*]," Published on Apr. 15, 2005 [retrieved on Mar. 27, 2002] [retrieved from the Internet:<URL:ncbi.nlm.nih.gov/protein/CAC24711>] [1 page].
- Genbank Accession No. U48435 [online], "*Solanum chacoense* putative cytochrome P450 gene, complete cds," Published on Jul. 2, 1997 [retrieved on Mar. 27, 2002] [retrieved from the Internet<URL:ncbi.nlm.nih.gov/protein/AAC39505>] [2 pages].
- Genbank Accession No. X96784 [online], "*N. tabacum* cytochrome P-450 gene," Published on Nov. 14, 2006 [retrieved on Mar. 27, 2002] [retrieved from the Internet:<URL:ncbi.nlm.nih.gov/nuccore/X96784>] [2 pages].
- Genbank Accession No. Y09447 [online], "*P. vulgaris* mRNA for cinnamate 4-hydroxylase," Published on Jul. 7, 1999 [retrieved on Mar. 27, 2002] [retrieved from the Internet:<URL:ncbi.nlm.nih.gov/nuccore/Y09447>] [1 page].
- Goeddel et al., "Synthesis of human fibroblast interferon by *E. coli*," *Nucleic Acids Res.* 8:4057-4074 (1980).
- Gonzalez et al., "Cytochromes P450 expression systems," *Annu. Rev. Pharmacol. Toxicol.* 35:369-390 (1995).
- Gordon-Kamm et al., "Transformation of maize cells and regeneration of fertile transgenic plants," *Plant Cell* 2:603-618 (1990).
- Gotoh, "Substrate recognition sites in cytochrome P450 family 2 (CYP2) proteins inferred from comparative analyses of amino acid and coding nucleotide sequences," *J. Biol. Chem.* 267:83-90 (1992).
- Greenhagen et al., "Identifying and manipulating structural determinates linking catalytic specificities in terpene synthases," *Proc. Natl. Acad. Sci. U.S.A.* 103:9826-9831 (2006).
- Greenhagen et al., "Probing sesquiterpene hydroxylase activities in a coupled assay with terpene synthases," *Arch. Biochem. Biophys.* 409:385-394 (2003).
- Greenhagen et al., "Molecular scaffolds for chemical wizardry: learning nature's rules for terpene cyclases," *Proc. Natl. Acad. Sci. U.S.A.* 98:13479-13491 (2001).
- Guarente et al., "A GAL 10-CYC1 hybrid yeast promoter identifies the GAL4 regulatory region as an upstream site," *Proc. Natl. Acad. Sci. U.S.A.* 79:7410-7414 (1982).
- Hahn, "Microbial elicitors and their receptors in plants," *Annu. Rev. Phytopathol.* 34:387-412 (1996).
- Hallahan et al., "Cytochrome-P450-catalysed monoterpenoid oxidation in catmint (*Nepeta racemosa*) and avocado (*Persea americana*); evidence for related enzymes with different activities," *Biochim. Biophys. Acta.* 1201:94-100 (1994).
- Hallahan et al., "Cytochrome P450 in plant/insect interactions: Geraniol 10-hydroxylase and the biosynthesis of iridoid monoterpenoids," *Drug Metabol. Drug Interact.* 12:369-382 (1995).
- Hanley et al., "Solubilization, partial purification, and immunodetection of squalene synthetase from tobacco cell suspension cultures," *Plant Physiol.* 98:215-220 (1992).

(56)

References Cited

OTHER PUBLICATIONS

- Haralampidis et al., "A new class of oxidosqualene cyclases directs synthesis of antimicrobial phytoprotectants in monocots," *Proc. Natl. Acad. Sci. U.S.A.* 98(23):13431-13436 (2001).
- Haralampidis et al., "Biosynthesis of triterpenoid saponins in plants," *Adv. Biochem. Eng. Biotechnol.* 75:31-49 (2002).
- Haudenschild et al., "Functional expression of regiospecific cytochrome P450 limonene hydroxylases from mint (*Mentha* spp.) in *Escherichia coli* and *Saccharomyces cerevisiae*," *Arch. Biochem. Biophys.* 379(1):127-136 (2000).
- Hefner et al., "Cytochrome P450-catalyzed hydroxylation of taxa-4(5),11(12)-diene to taxa-4(20),11(12)-dien-5 α -ol: the first oxygenation step in taxol biosynthesis," *Chem. Biol.* 3(6):479-489 (1996).
- Helliwell et al., "*Arabidopsis* ent-kaurene oxidase catalyzes three steps of gibberellin biosynthesis," *Plant Physiol.* 119:507-510 (1999).
- Helliwell et al., "Cloning of the *Arabidopsis* ent-kaurene oxidase gene GA3," *Proc. Natl. Acad. Sci. U.S.A.* 95:9019-9024 (1998).
- Holton et al., "Cloning and expression of cytochrome P450 genes controlling flower colour," *Nature* 366:276-279 (1993).
- Hoshino et al., "5-*epi*-Aristolochene 3-hydroxylase from green pepper," *Phytochemistry* 38:609-613 (1995).
- Humphreys et al., "Molecular 'phanning' with plant P450s," *Trends in Plant Science* 5:271-272 (2000).
- Hutvagner et al., "Cytochrome P450 71D6," UniProtKB/Swiss-Prot entry P93530, updated last on Aug. 10, 2010, Version 53, <URL: uniprot.org/uniprot/P93530 [accessed on Nov. 1, 2010] [5 pages].
- Hutvagner et al., "Cytochrome P450 71D7," UniProtKB/Swiss-Prot entry P93531, updated last on Aug. 10, 2010, Version 53, <URL: uniprot.org/uniprot/P93531 [accessed on Nov. 1, 2010] [5 pages].
- Hutvagner et al., "Isolation and sequence analysis of a cDNA and a related gene for cytochrome P450 proteins from *Solanum chacoense*," *Gene* 188:247-252 (1998).
- Irmeler et al., "Indole alkaloid biosynthesis in *Catharanthus roseus*: new enzyme activities and identification of cytochrome P450 CYP72A1 as secologanin synthase," *Plant J.* 24(6):797-804 (2000).
- Jang et al., "Sugar sensing in higher plants," *Plant Cell* 6:1665-1679 (1994).
- Kay et al., "Duplication of CaMV 35S promoter sequences creates a strong enhancer for plant genes," *Science* 236:1299-1302 (1987).
- Keller et al., "Sesquiterpene cyclase is not a determining factor for elicitor- and pathogen-induced capsidiol accumulation in tobacco," *Planta* 205:467-476 (1998).
- Kindle, "High-frequency nuclear transformation of *Chlamydomonas reinhardtii*," *Proc. Natl. Acad. Sci. U.S.A.* 87:1228-1232 (1990).
- Koepf et al., "Cyclization of geranylgeranyl diphosphate to taxa-4(5),11(12)-diene is the committed step of taxol biosynthesis in Pacific Yew," *J. Biol. Chem.* 270:8686-8690 (1995).
- Kuhlemeier et al., "The pea rbcS-3A promoter mediates light responsiveness but not organ specificity," *Plant Cell* 1:471-478 (1989).
- Lupien et al., "Regiospecific cytochrome P450 limonene hydroxylases from mint (*Mentha*) species: cDNA isolation, characterization, and functional expression of (-)-4S-limonene-3-hydroxylase and (-)-4S-limonene-6-hydroxylase," *Arch. Biochem. Biophys.* 368:181-192 (1999).
- Mandujano-Chavez et al., "Differential induction of sesquiterpene metabolism in tobacco cell suspension cultures by methyl jasmonate and fungal elicitor," *Arch. Biochem. Biophys.* 381:285-294 (2000).
- Marcotte et al., "Absciscic acid-responsive sequences from the Em gene of wheat," *Plant Cell* 1:969-976 (1989).
- Mathis et al., "Pre-steady-state study of recombinant sesquiterpene cyclases," *Biochem. J.* 36:8340-8348 (1997).
- Maughan et al., "Expression of CYP71B7, a cytochrome P450 expressed sequence tag from *Arabidopsis thaliana*," *Arch. Biochem. Biophys.* 341:104-111 (1997).
- Milet et al., "Capsidiol and ethylene production by tobacco cells in response to cryptogin, an elicitor from *Phytophthora cryptogea*," *Phytochemistry* 30:2171-2173 (1991).
- Miller, "Structure of genes encoding steriodogenic enzymes," *J. Steroid. Biochem.* 27:759-766 (1987).
- Molot et al., "Relations between capsidiol concentration, speed of fungal invasion and level of induced resistance in cultivars of pepper (*Capsicum annuum*) susceptible or resistant to *Phytophthora capsici*," *Physiol. Plant Pathol.* 18:379-389 (1981).
- Nedelkina et al., "Novel characteristics and regulation of a divergent cinnamate 4-hydroxylase (CYP3A15) from french bean: engineering expression in yeast," *Plant Mol. Biol.* 39:1079-1090 (1999).
- Nelson et al., "Evolution of cytochrome P-450 proteins," *Mol. Biol. Evol.* 4(6):572-593 (1987).
- Newman et al., "Characterization of the TAC box, a cis-element within an elicitor-inducible sesquiterpene cyclase promoter," *Plant J.* 16:1-12 (1998).
- Nomura et al., "The cDNA cloning and transient expression of a chicken gene encoding cytochrome P-450scc," *Gene* 185(2):217-222 (1997).
- Nunez et al., "Isolation of the putative cDNA encoding cholesterol side chain cleavage cytochrome P450 (CYP11A) of the southern stingray (*Dasyatis americana*)," *Gene* 187(1):123-129 (1997).
- Odell et al., "Identification of DNA sequences required for activity of the cauliflower mosaic virus 25S promoter," *Nature* 313:810-812 (1985).
- O'Donohue et al., "Chemical synthesis, expression and mutagenesis of a gene encoding beta-cryptogin, an elicitor produced by *Phytophthora cryptogea*," *Plant Mol. Biol.* 27:577-586 (1995).
- Ohnuma et al., "A role of the amino acid residue located on the fifth position before the first aspartate-rich motif of farnesyl diphosphate synthase of determination of the final product," *J. Biol. Chem.* 271:30748-30754 (1996).
- O'Keefe et al., "Cytochrome P-450 from the mesocarp of avocado (*Persea americana*)," *Plant Physiol.* 89:1141-1149 (1989).
- O'Maille et al., "Biosynthetic potential of sesquiterpene synthases: alternative products of tobacco 5-*epi*-aristolochene synthase," *Arch. Biochem. Biophys.* 448:73-82 (2006).
- Omura et al., "The carbon monoxide-binding pigment of liver microsomes I. Evidence for its hemoprotein nature," *J. Biol. Chem.* 239:2370-2378 (1964).
- Omura, "Forty years of cytochrome P450," *Biochem. Biophys. Res. Commun.* 266:690-698 (1999).
- Ow et al., "Functional regions of the cauliflower mosaic virus 35S RNA promoter determined by use of the Firefly luciferase gene as a reporter of promoter activity," *Proc. Natl. Acad. Sci. U.S.A.* 84:4870-4874 (1987).
- Pompon et al., "Yeast expression of animal and plant P450s in optimized redox environments," *Methods Enzymol.* 272:51-64 (1996).
- Porter et al., "Strategies to enhance the coexpression of cytochrome P450 2E1 and reductase in bacteria," *Drug Metab. Rev.* 31:159-174 (1999).
- Potrykus, "Gene transfer to plants: assessment of published approaches and results," *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 42:205-225 (1991).
- Rademacher, "Growth retardants: effects on gibberellin biosynthesis and other metabolic pathways," *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 51:501-531 (2000).
- Ralston et al., "Biochemical and molecular characterization of 5-*epi*-aristolochene 3-hydroxylase, a putative regulatory enzyme in the biosynthesis of sesquiterpene phytoalexins in tobacco," *Plant Interactions with Other Organisms. Annual Meeting of the American Society of Plant Physiologists. Madison, WI., Jun. 27-Jul. 1, 1998, Session 47: Abstract #737 (Poster Presentation).*
- Ralston et al., "Cloning, heterologous expression, and functional characterization of 5-*epi*-aristolochene-1,3-dihydroxylase from tobacco (*Nicotiana tabacum*)," *Arch. Biochem. Biophys.* 393:222-235 (2001).
- Rising et al., "Demonstration of gerrnacrene A as an intermediate in 5-*epi*-aristolochene synthesis catalysis," *J. Am. Chem. Soc.* 122:1861-1866 (2000).
- Rosahl et al., "Expression of a tuber-specific storage protein in transgenic tobacco plants: demonstration of an esterase activity," *EMBO J.* 6:1155-1159 (1987).

(56)

References Cited

OTHER PUBLICATIONS

- Schäffner et al., "Maize *rbcs* promoter activity depends on sequence elements not found in dicot *rbcs* promoters," *Plant Cell* 3:997-1012 (1991).
- Schalk et al., "A single amino acid substitution (F363I) converts the regiochemistry of the spearmint (-)-limonene hydroxylase from a C6- to a C3-hydroxylase," *Proc. Natl. Acad. Sci. U.S.A.* 97:11948-11953 (2000).
- Schenk et al., "Stereochemistry and deuterium isotope effects associated with the cyclization-rearrangements catalyzed by tobacco epiaristolochene and *hyoscyamus* premnaspirodiene synthases, and the chimeric CH4 hybrid cyclase," *Arch. Biochem. Biophys.* 448:31-44 (2006).
- Scherthaner et al., "Endosperm-specific activity of a zein gene promoter in transgenic tobacco plants," *EMBO J.* 7:1249-1255 (1988).
- Schoendorf et al., "Molecular cloning of a cytochrome P450 taxane 10 β -hydroxylase cDNA from *Taxus* and functional expression in yeast," *Proc. Natl. Acad. Sci. U.S.A.* 98(4):1501-1506 (2001).
- Schopfer et al., "Identification of elicitor-induced cytochrome P450s of soybean (*Glycine max* L.) using differential display of mRNA," *Mol. Gen. Genet.* 258(4):315-322 (1998).
- Schuler, "Plant cytochrome P450 monooxygenases," *Crit. Rev. Plant Sci.* 15:235-284 (1996).
- Sheen et al., "Green-fluorescent protein as a new vital marker in plant cells," *Plant J.* 8:777-784 (1995).
- Sheen, "Metabolic repression of transcription in higher plants," *Plant Cell* 2:1027-1038 (1990).
- Shen et al., "Functional dissection of an abscisic acid (ABA)-inducible gene reveals two independent ABA-responsive complexes each containing a G-box and a novel cis-acting element," *Plant Cell* 7:295-307 (1995).
- Shimatake et al., "Purified λ regulatory protein cII positively activates promoters for lysogenic development," *Nature* 292:128-132 (1981).
- Siebertz et al., "cis-Analysis of the wound-inducible promoter *wun* 1 in transgenic tobacco plants and histochemical localization of its expression," *Plant Cell* 1:961-968 (1989).
- Simpson et al., "Light-inducible and tissue-specific expression of a chimeric gene under control of the 5'-flanking sequence of a pea chlorophyll *a/b*-binding protein gene," *EMBO J.* 4:2723-2729 (1985).
- Starks et al., "Structural basis for cyclic terpene biosynthesis by tobacco 5-*epi*-aristolochene synthase," *Science* 277:1815-1820 (1997).
- Stolle et al., "Restricted colonization by *Peronospora tabacina* and phytoalexin accumulation in immunized tobacco leaves," *Phytopathology* 78:1193-1197 (1988).
- Straub et al., "Structure and promoter analysis of an ABA- and stress-regulated barley gene, *HVA1*," *Plant Mol. Biol.* 6:617-630 (1994).
- Sutherland et al., "A cytochrome P450 terpenoid hydroxylase linked to the suppression of insect juvenile hormone synthesis," *Proc. Natl. Acad. Sci. U.S.A.* 95:12884-12889 (1998).
- Takahashi et al., "Functional characterization of premnaspirodiene oxygenase, a cytochrome P450 catalyzing regio- and stereo-specific hydroxylations of diverse sesquiterpene substrates," *J. Biol. Chem.* 278:31744-31754 (2003).
- Takahashi et al., "Kinetic and molecular analysis of 5-*epi*-aristolochene 1,3-dihydroxylase, a cytochrome P450 enzyme catalyzing successive hydroxylations of sesquiterpenes," *J. Biol. Chem.* 280:3686-3696 (2005).
- Takahashi et al., "Metabolic engineering of sesquiterpene metabolism in yeast," *Biotechnol. Bioeng.* 97:170-181 (2007).
- Takahashi et al., "Characterization of two genes encoding small heat-shock proteins in *Arabidopsis thaliana*," *Mol. Gen. Genet.* 219:365-372 (1989).
- Takemoto et al., "Molecular cloning of a defense-response-related cytochrome P450 gene from tobacco," *Plant Cell Physiol.* 40:1232-1242 (1999).
- Tarshis et al., "Regulation of product chain length by isoprenyl diphosphate synthases," *Proc. Natl. Acad. Sci. U.S.A.* 93:15018-15023 (1996).
- Terada et al., "Expression of CaMV35S-GUS gene in transgenic rice plants," *Mol. Gen. Genet.* 220:389-392 (1990).
- Thai et al., "Farnesol is utilized for isoprenoid biosynthesis in plant cells via farnesyl pyrophosphate formed by successive monophosphorylation reactions," *Proc. Natl. Acad. Sci. U.S.A.* 96:13080-13085 (1999).
- Thornburg et al., "Wound-inducible expression of a potato inhibitor II-chloramphenicol acetyltransferase gene fusion in transgenic tobacco plants," *Proc. Natl. Acad. Sci. U.S.A.* 84:744-748 (1987).
- Threlfall et al., "Co-ordinated inhibition of squalene synthetase and induction of enzymes of sesquiterpenoid phytoalexin biosynthesis in cultures of *Nicotiana tabacum*," *Phytochem.* 27:2567-2580 (1988).
- Trant, "Isolation and characterization of the cDNA encoding the spiny dogfish shark (*Squalus acanthias*) form of cytochrome P450c17," *J. Exp. Zool.* 272(1):25-33 (1995).
- Trant, "Functional expression of recombinant spiny dogfish shark (*Squalus acanthias*) cytochrome P450c17 (17 α -hydroxylase/C17,20-lyase) in yeast (*Pichia pastoris*)," *Arch. Biochem. Biophys.* 326(1):8-14 (1996).
- Tudzynski, "Biosynthesis of gibberellins in *Gibberella fujikuroi*: biomolecular aspects," *Appl. Microbiol. Biotechnol.* 52(3):298-310 (1999).
- Umemoto et al., "cDNAs sequences encoding cytochrome P450 (CYP71 family) from eggplant seedlings," *FEBS* 330(2):169-173 (1993).
- Urban et al., "Cloning, yeast expression, and characterization of the coupling of two distantly related *Arabidopsis thaliana* NADPH-cytochrome P450 reductases with P450 CYP73A5," *J. Biol. Chem.* 272:19176-19186 (1997).
- Urban et al., "Maximizing the expression of mammalian cytochrome P-450 monooxygenase activities in yeast cells," *Biochimie* 72:463-472 (1990).
- Vögeli et al., "Inhibition of phytoalexin biosynthesis in elicitor-treated tobacco cell-suspension cultures by calcium/calmodulin antagonists," *Plant Physiol.* 100:1369-1376 (1992).
- Vögeli et al., "Purification and characterization of an inducible sesquiterpene cyclase from elicitor-treated tobacco cell suspension cultures," *Plant Physiol.* 93:182-187 (1990).
- Vögeli et al., "Induction of sesquiterpene cyclase and suppression of squalene synthetase activities in plant cell cultures treated with fungal elicitor," *Plant Physiol.* 88:1291-1296 (1988).
- Vögeli et al., "Inhibition of a plant sesquiterpene cyclase by mevinolin," *Arch. Biochem. Biophys.* 288:157-162 (1991).
- Vögeli et al., "Regulation of a sesquiterpene cyclase in callus-treated tobacco cell suspension cultures," *Plant Physiol.* 94:1860-1866 (1990).
- Walker et al., "Molecular cloning of a 10-deacetylbaicatin III-10-O-acetyl transferase cDNA from *Taxus* and functional expression in *Escherichia coli*," *Proc. Natl. Acad. Sci. U.S.A.* 97:583-587 (2000).
- Watson et al., "Sesquiterpenoid stress metabolites in capsicums," *Biochem. Soc. Trans.* 11:589 (1983).
- Werck-Reichhart et al., "Cytochromes P450 for engineering herbicide tolerance," *Trends in Plant Science* 5:116-123 (2000).
- Werck-Reichhart et al., "Cytochromes P450: a success story," *Genome Biol.* 1(6)Reviews:3003.1-3003.9 (2000).
- Whitehead et al., "5-*epi*-Aristolochene is a common precursor of the sesquiterpene phytoalexins capsidiol and debneyol," *Phytochemistry* 28:775-779 (1989).
- Whitehead et al., "Synthesis of (+)-5-*epi*-aristolochene and (+)-1-deoxycapsidiol from capsidiol," *Phytochemistry* 29:479-482 (1990).
- Whitehead, "Cis-9,10-dihydrocapsenone: a possible catabolite of capsidiol from cell suspension cultures of *Capsicum annuum*," *Phytochemistry* 26:1367-1369 (1987).
- Wildung et al., "A cDNA clone for taxadiene synthase, the diterpene cyclase that catalyzes the committed step of taxol biosynthesis," *J. Biol. Chem.* 271:9201-9204 (1996).
- Williams et al., "Intramolecular proton transfer in the cyclization of geranylgeranyl diphosphate to the taxadiene precursor of taxol catalyzed by recombinant taxadiene synthase," *Chem. Biol.* 7:969-977 (2000).

(56)

References Cited

OTHER PUBLICATIONS

Wu et al., "Expression cloning and characterization of human 17 beta-hydroxysteroid dehydrogenase type 2, a microsomal enzyme possessing 20 alpha-hydroxysteroid dehydrogenase activity," J. Biol. Chem. 268(17):12964-12969 (1993).

Wu et al., "Redirection of cytosolic or plastidic isoprenoid precursors elevates terpene production in plants," Nat. Biotechnol. 24:1441-1447 (2006).

Wu et al., "Surrogate splicing for functional analysis of sesquiterpene synthase genes," Plant Physiol. 138:1322-1333 (2005).

Wust et al., "Hydroxylation of limonene enantiomers and analogs by recombinant (-)-limonene 3- and 6-hydroxylases from mint (*Mentha*) species: evidence for catalysis within sterically constrained active sites," Arch. Biochem. Biophys. 387:125-136 (2001).

Yin et al., "Regulation of sesquiterpene cyclase gene expression—characterization of an elicitor- and pathogen-inducible promoter," Plant Physiol. 115:437-451 (1997).

Zhang et al., "Efficient regeneration of transgenic plants from rice protoplasts and correctly regulated expression of the foreign gene in the plants," Theor. Appl. Genet. 76:835-840 (1988).

Zhao et al., "Eremophilane sesquiterpenes from capsidiol," J. Org. Chem. 69:7428-7435 (2004).

Zook et al., "Characterization of novel sesquiterpene biosynthesis in tobacco expressing fungal sesquiterpenoid synthase," Plant Physiol. 112:311-318 (1996).

Letter/Written Disclosure of the Supplemental Information Disclosure Statement for the above-referenced application, mailed on the same day herewith, 2 pages.

Examination Report, issued Jun. 11, 2013, in connection with corresponding Australian Patent Application No. 2012202780, 2 pages.

Notice of Acceptance, issued Jun. 24, 2013, in connection with corresponding Australian Patent Application No. 2013204829, 2 pages. US 8,486,659, 07/2013, Julien et al. (withdrawn)

* cited by examiner

FIG. 1




FIG. 2




FIG. 3A




FIG. 3B




FIG. 4A



FIG. 4B

KETLRLH-for	5'-AARGARACIYTIMGIYTICA-3'
KETLRLY-for	5'-AARGARACIYTIMGIYTITA-3'
KETLRLR-for	5'-AARGARACIYTIMGIYTIMG-3'
FXPERF -for	5'-TTYIIICCIGARMGITY-3'
FXPERF-rev	5'-RAAICKYTCIGGIIIRAA-3'
GRRXCP(A/G)-for	5'-GGIMGIMGIIIIITGYCCIGS-3'
PFGXGRR-rev	5'-CKICKICCIIIICCRAAIGG-3'
T7	5'-GTAATACGACTCACTATAGGG-3'
T3	5'-CAATTAACCCTCACTAAAGGG-3'

FIG. 4C




FIG. 5




FIG. 6A






FIG. 6B





MpCYP71D13	33	E	P	Q	-	W	R	K	K	P	-	K	P	Q	-	E
MsCYP71D18	32	M	E	L	D	L	L	S	A	I	I	L	L	S	A	I
NrCYP71A5v1	29	M	A	I	L	S	A	I	I	I	I	L	L	S	A	I
PaCYP71A1	33	M	A	I	L	S	A	I	I	I	I	L	L	S	A	I
NiCYP71D20	30	M	Q	F	F	S	L	V	S	L	L	F	L	F	L	S
AtCYP701A3	39	M	A	F	F	S	M	I	S	I	L	L	G	F	V	I
MpCYP71D13	70	N	L	P	P	G	P	-	-	P	K	L	P	P	L	P
MsCYP71D18	69	N	L	P	P	S	P	-	-	P	K	L	P	P	V	V
NrCYP71A5v1	67	N	L	P	P	S	P	-	-	P	K	L	P	P	V	V
PaCYP71A1	65	N	L	P	P	S	P	-	-	P	K	L	P	P	V	V
NiCYP71D20	67	K	L	P	P	G	P	-	-	W	K	I	P	P	I	P
AtCYP701A3	78	T	L	P	S	V	P	V	P	G	F	P	V	P	G	F
MpCYP71D13	109	A	H	V	Q	L	G	E	V	F	S	V	V	V	L	L
MsCYP71D18	108	A	H	V	Q	L	G	E	V	F	S	V	V	V	L	L
NrCYP71A5v1	106	M	Q	L	H	F	G	S	V	P	T	L	I	V	V	A
PaCYP71A1	104	I	L	L	H	L	G	H	I	P	T	L	I	V	V	A
NiCYP71D20	106	M	H	L	Q	L	G	E	I	S	A	V	V	V	L	L
AtCYP701A3	117	Y	S	I	K	M	G	S	S	L	I	V	V	L	L	S
MpCYP71D13	147	K	I	M	W	Y	D	N	D	D	D	D	D	D	D	D
MsCYP71D18	146	R	T	M	W	Y	D	N	D	D	D	D	D	D	D	D
NrCYP71A5v1	144	N	R	L	F	F	N	N	R	R	T	D	D	D	D	D
PaCYP71A1	142	R	R	I	F	F	N	N	R	R	T	D	D	D	D	D
NiCYP71D20	144	D	I	C	Y	N	C	S	S	M	V	A	T	S	D	Y
AtCYP701A3	155	T	V	L	T	C	D	K	S	M	V	A	T	S	D	Y

Figure 8A

[illegible]

Figure 8B

MpCYP71D13	I	T	T	N	A	I	K	A	F	I	F	D	T	F	S	A	G	T	E	T	S	S	T	T	L	T	T	W	V	M	A	E	L	M	R	N	P	E	V	M	321		
MsCYP71D18	I	T	S	N	C	I	K	K	G	F	I	D	T	F	S	A	G	A	E	T	S	S	T	T	I	S	T	W	A	L	S	E	L	M	R	N	P	A	K	M	321		
NrCYP71A5v1	V	E	D	N	T	V	K	K	A	L	L	D	M	F	F	A	G	T	D	T	S	S	T	A	L	E	W	A	M	A	E	L	I	K	N	P	R	A	M	332			
PaCYP71A1	L	N	R	N	N	L	K	K	A	V	L	D	M	F	F	A	G	T	D	T	S	S	T	V	E	W	A	M	A	E	L	M	K	N	P	D	V	M	327				
NiCYP71D20	-	-	-	-	-	-	-	-	-	G	A	L	E	T	I	E	T	A	D	T	S	S	T	L	T	T	T	W	A	M	A	E	L	M	K	N	P	S	V	F	295		
AtCYP701A3	L	T	K	E	Q	I	A	I	L	V	I	D	T	F	S	A	G	T	E	T	S	S	T	L	T	T	T	W	A	A	Y	E	L	A	K	N	P	S	V	Q	339		
MpCYP71D13	A	K	A	Q	A	E	V	R	A	A	L	K	G	K	T	D	V	D	V	D	S	D	V	Q	E	L	K	Y	M	K	S	V	V	K	E	T	M	R	M	H	363		
MsCYP71D18	A	K	V	Q	A	E	V	R	E	A	L	K	G	S	K	A	V	E	I	E	E	D	L	E	K	P	Y	L	R	S	S	I	V	V	K	E	T	L	R	L	H	360	
NrCYP71A5v1	K	R	L	Q	N	Q	E	V	R	E	V	A	G	S	K	A	V	S	F	D	E	E	D	L	E	K	P	Y	L	R	S	S	I	V	V	K	E	T	L	R	L	H	371
PaCYP71A1	E	K	A	Q	A	E	V	R	E	V	V	A	G	S	K	A	V	S	F	D	E	E	D	L	E	K	P	Y	L	R	S	S	I	V	V	K	E	T	L	R	L	H	366
NiCYP71D20	T	K	A	Q	A	E	V	R	E	V	A	F	C	G	E	-	K	F	K	E	E	Q	L	S	Q	V	Y	L	R	S	S	I	V	V	K	E	T	L	R	L	H	334	
AtCYP701A3	D	R	L	C	K	E	I	Q	N	V	C	G	G	E	-	K	F	K	E	E	Q	L	S	Q	V	L	Y	L	R	S	S	I	V	V	K	E	T	L	R	K	Y	377	
MpCYP71D13	P	P	F	P	-	-	L	I	P	-	-	R	S	Q	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	400
MsCYP71D18	P	P	F	P	-	-	L	I	P	-	-	R	S	Q	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	397
NrCYP71A5v1	V	P	V	V	L	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	409
PaCYP71A1	P	V	A	P	L	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	404
NiCYP71D20	P	S	A	P	L	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	372
AtCYP701A3	S	P	A	P	-	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	415
MpCYP71D13	P	P	F	P	-	-	L	I	P	-	-	R	S	Q	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	439
MsCYP71D18	P	P	F	P	-	-	L	I	P	-	-	R	S	Q	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	436
NrCYP71A5v1	V	P	V	V	L	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	448
PaCYP71A1	P	V	A	P	L	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	443
NiCYP71D20	P	S	A	P	L	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	411
AtCYP701A3	S	P	A	P	-	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	454
MpCYP71D13	P	P	F	P	-	-	L	I	P	-	-	R	S	Q	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	439
MsCYP71D18	P	P	F	P	-	-	L	I	P	-	-	R	S	Q	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	436
NrCYP71A5v1	V	P	V	V	L	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	448
PaCYP71A1	P	V	A	P	L	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	443
NiCYP71D20	P	S	A	P	L	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	411
AtCYP701A3	S	P	A	P	-	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	454
MpCYP71D13	P	P	F	P	-	-	L	I	P	-	-	R	S	Q	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	439
MsCYP71D18	P	P	F	P	-	-	L	I	P	-	-	R	S	Q	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	436
NrCYP71A5v1	V	P	V	V	L	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	448
PaCYP71A1	P	V	A	P	L	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	443
NiCYP71D20	P	S	A	P	L	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	411
AtCYP701A3	S	P	A	P	-	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	454
MpCYP71D13	P	P	F	P	-	-	L	I	P	-	-	R	S	Q	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	439
MsCYP71D18	P	P	F	P	-	-	L	I	P	-	-	R	S	Q	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	436
NrCYP71A5v1	V	P	V	V	L	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	448
PaCYP71A1	P	V	A	P	L	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	443
NiCYP71D20	P	S	A	P	L	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	411
AtCYP701A3	S	P	A	P	-	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	454
MpCYP71D13	P	P	F	P	-	-	L	I	P	-	-	R	S	Q	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	439
MsCYP71D18	P	P	F	P	-	-	L	I	P	-	-	R	S	Q	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	436
NrCYP71A5v1	V	P	V	V	L	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	448
PaCYP71A1	P	V	A	P	L	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	443
NiCYP71D20	P	S	A	P	L	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	411
AtCYP701A3	S	P	A	P	-	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	454
MpCYP71D13	P	P	F	P	-	-	L	I	P	-	-	R	S	Q	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	439
MsCYP71D18	P	P	F	P	-	-	L	I	P	-	-	R	S	Q	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	436
NrCYP71A5v1	V	P	V	V	L	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	448
PaCYP71A1	P	V	A	P	L	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R	I	M	F	L	L	I	N	N	A	443
NiCYP71D20	P	S	A	P	L	-	L	V	P	-	-	R	E	E	S	T	R	D	E	E	C	E	V	N	G	Y	T	I	P	N	K	A	R										

Figure 8C

Figure 8D

CYTOCHROME P450S AND USES THEREOF

RELATED APPLICATIONS

This application is a continuation of U.S. patent application Ser. No. 13/199,349, filed Aug. 26, 2011 (now issued U.S. Pat. No. 8,445,231), which is a continuation of U.S. patent application Ser. No. 12/182,000, filed Jul. 29, 2008 (now issued U.S. Pat. No. 8,263,362), which is a continuation of U.S. patent application Ser. No. 10/097,559, filed Mar. 8, 2002 (now issued U.S. Pat. No. 7,405,057), which claims the benefit of U.S. Provisional Application Nos. 60/274,421 and 60/275,597, filed on Mar. 9, 2001 and Mar. 13, 2001, respectively, all of which are hereby incorporated by reference.

FIELD OF THE INVENTION

This invention relates to cytochrome P450s and uses thereof.

BACKGROUND OF THE INVENTION

Cytochrome P450s encompass a superfamily of oxidases responsible for the oxidation of numerous endobiotics and thousands of xenobiotics. In addition, in plants, cytochrome P450s play important roles in wound healing, pest resistance, signaling, and anti-microbial and anti-fungal activity.

Capsidiol is a bicyclic, dihydroxylated sesquiterpene produced by many Solanaceous species in response to a variety of environmental stimuli, including exposure to UV (Back et al., *Plant Cell. Physiol.* 389:899-904, 1998) and infection by microorganisms (Molot et al., *Physiol. Plant Pathol.* 379-389, 1981; Stolle et al., *Phytopathology* 78:1193-1197, 1988; Keller et al., *Planta*. 205:467-476, 1998). It is the primary antibiotic or phytoalexin produced in tobacco in response to fungal elicitation, and it is derived from the isoprenoid pathway via its hydrocarbon precursor, 5-epi-aristolochene (FIG. 1). Several of the biosynthetic enzymes leading up to 5-epi-aristolochene formation have been studied (Chappell, *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 46:521-547, 1995), especially 5-epi-aristolochene synthase (BAS) (Vogeli and Chappell, *Plant Physiol.* 88:1291-1296, 1988; Back and Chappell, *Proc. Natl. Acad. Sci. U.S.A.* 93:6841-6845, 1996; Mathis et al., *Biochemistry* 36:8340-8348, 1997; Starks et al., *Science* 277: 1815-1820, 1997). BAS commits carbon to sesquiterpene metabolism by catalyzing the cyclization of farnesyl diphosphate (FPP) to 5-epi-aristolochene. However, until the present invention, the enzyme(s) responsible for the conversion of 5-epi-aristolochene to capsidiol has yet to be fully identified and characterized.

Biochemical evidence from previous studies in tobacco (Whitehead et al., *Phytochemistry* 28:775-779, 1989) and green pepper (Hoshino et al., *Phytochemistry* 38:609-613, 1995) have suggested that the oxidation of 5-epi-aristolochene to capsidiol occurs in a two step process with one of the hydroxylation steps being constitutive and the other being mediated by an elicitor-inducible cytochrome P450 (FIG. 1). Because 1-deoxycapsidiol had been isolated from natural sources (Watson et al., *Biochem. Soc. Trans.* 11:589, 1983), Whitehead et al. (*Phytochemistry* 28:775-779, 1989), surmised that perhaps the biosynthesis of this intermediate was due to pathogen induction of a corresponding hydroxylase. They therefore prepared synthetic 1-deoxycapsidiol and reported a modest conversion of this compound to capsidiol when fed to control or unelicited tobacco cell cultures. This was further supported by their observation that radiolabeled 5-epi-aristolochene was only converted to capsidiol when fed

to elicitor-induced cell cultures but not control cultures. Whitehead et al. (*Phytochemistry* 28:775-779, 1989) therefore concluded that the 3-hydroxylase, responsible for hydroxylation of 5-epi-aristolochene at C3 to generate 1-deoxycapsidiol, was pathogen/elicitor inducible, while the 1-hydroxylase, responsible for hydroxylating 1-deoxycapsidiol at the C1 to generate capsidiol, was constitutive. Hoshino et al. (*Phytochemistry* 38:609-613, 1995) added to the observations of Whitehead et al. (*Phytochemistry* 28:775-779, 1989) by directly measuring 3-hydroxylase-activity in microsomal preparations of arachidonic acid-elicited *Capsicum annuum* fruits and seedlings. These assays consisted of incubating 5-epi-aristolochene with microsome preparations and subsequently determining the amount of 1-deoxycapsidiol generated by a combination of thin-layer chromatography (TLC) separations and gas chromatography (GC). Their evidence demonstrated that the conversion of 5-epi-aristolochene to 1-deoxycapsidiol was dependent on both NADPH and O₂, and that 1-deoxycapsidiol accumulation in vitro was arrested by the P450 antagonists carbon monoxide (Omura and Sato, *J. Biol. Chem.* 239:2370-2378, 1964), ancyimidol (Coolbaugh et al., *Plant Physiol.* 62:571-576, 1978), and ketoconazole (Rademacher, *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 51:501-531, 2000).

Recent results suggest that the hydroxylation of 5-epi-aristolochene is an important regulated step in capsidiol biosynthesis. In studies to evaluate the effectiveness of methyl-jasmonate as an inducer of capsidiol biosynthesis in tobacco cell cultures, Mandujano-Chávez et al. (*Arch. Biochem. Biophys.* 381:285-294, 2000), reported that the modest accumulation of this phytoalexin was accompanied by a strong induction of EAS. This result implied that steps before or after the sesquiterpene cyclase reaction were limiting. Using an in vivo assay measuring the conversion rate of radiolabeled 5-epi-aristolochene to capsidiol, a very limited induction of the hydroxylase activity was observed in cells treated with methyl jasmonate relative to that in fungal elicitor-treated cells. This result pointed to the hydroxylase reactions as a potentially limiting step in capsidiol biosynthesis.

SUMMARY OF THE INVENTION

In one aspect, the invention features several isolated cytochrome P450 polypeptides (such as CYP71D20, CYP71D21, CYP73A27, CYP73A28, and CYP92A5, and P450s having substantial identity to these polypeptides), as well as isolated nucleic acid molecules that encode these P450s.

In related aspects, the invention features a vector (such as an expression vector) including an isolated nucleic acid molecule of the invention and a cell (for example, a prokaryotic cell, such as *Agrobacterium* or *E. coli*, or a eukaryotic cell, such as a mammalian, insect, yeast, or plant cell) including the isolated nucleic acid molecule or vector.

In yet another aspect, the invention features a transgenic plant or transgenic plant component including a nucleic acid molecule of the invention, wherein the nucleic acid molecule is expressed in the transgenic plant or the transgenic plant component. Preferably, the transgenic plant or transgenic plant component is an angiosperm (for example, a monocot or dicot). In preferred embodiments, the transgenic plant or transgenic plant component is a solanaceous, maize, rice, or cruciferous plant or a component thereof. The invention further includes a seed produced by the transgenic plant or transgenic plant component, or progeny thereof.

In another aspect, the invention features a method of providing an increased level of resistance against a disease caused by a plant pathogen in a transgenic plant. The method

involves: (a) producing a transgenic plant cell including the nucleic acid molecule of the invention integrated into the genome of the transgenic plant cell and positioned for expression in the plant cell; and (b) growing a transgenic plant from the plant cell wherein the nucleic acid molecule is expressed in the transgenic plant and the transgenic plant is thereby provided with an increased level of resistance against a disease caused by a plant pathogen.

In another aspect, the invention features a method for producing an altered compound, the method including the steps of contacting the compound with one or more of the isolated polypeptides disclosed herein under conditions allowing for the hydroxylation, oxidation, demethylation, or methylation of the compound and recovering the altered compound.

In still another aspect, the invention features a hydroxylating agent including any of the isolated polypeptides disclosed herein.

In yet another embodiment, the invention features an isolated nucleic acid molecule that specifically hybridizes under highly stringent conditions to the complement of any one of the sequences described in SEQ ID NO:2 (CYP71D20), SEQ ID NO:4 (CYP71D21), SEQ ID NO:6 (CYP73A27), SEQ ID NO:8 (CYP73A28), or SEQ ID NO:12 (CYP92A5), wherein such a nucleic acid molecule encodes a cytochrome P450 polypeptide.

In another aspect, the invention features a host cell expressing a recombinant isoprenoid synthase and a recombinant cytochrome P450. In preferred embodiments, the host cell further expresses, independently or in combination, a recombinant acetyltransferase, methyltransferase, or fatty acyltransferase. In other preferred embodiments, the host expresses an endogenous or recombinant cytochrome reductase. Preferably, the host cell is a yeast cell, a bacterial cell, an insect cell, or a plant cell.

In a related aspect, the invention features a method for producing an isoprenoid compound, the method including the steps of: (a) culturing a cell that expresses a recombinant isoprenoid synthase and a recombinant cytochrome P450 under conditions wherein the isoprenoid synthase and the cytochrome P450 are expressed and catalyze the formation of an isoprenoid compound not normally produced by the cell; and (b) recovering the isoprenoid compound. In preferred embodiments, the host cell further expresses a recombinant acetyltransferase, a recombinant methyltransferase, or a recombinant fatty acyltransferase. In other preferred embodiments, the host cell expresses an endogenous or recombinant cytochrome reductase. Preferably, the host cell is a yeast cell, a bacterial cell, an insect cell, or a plant cell.

In yet another aspect, the invention features an isoprenoid compound produced according to the above-mentioned methods.

By "P450 polypeptide," "cytochrome P450," or "P450" is meant a polypeptide that contains a heme-binding domain and shows a CO absorption spectra peak at 450 nm according to standard methods, for example, those described herein. Such P450s may also include, without limitation, hydroxylase activity, dual hydroxylase activity, demethylase activity, or oxidase activity. Such enzymatic activities are determined using methods well known in the art.

By "polypeptide" is meant any chain of amino acids, regardless of length or post-translational modification (for example, glycosylation or phosphorylation).

By "substantially identical" is meant a polypeptide or nucleic acid exhibiting at least 80 or 85%, preferably 90%, more preferably 95%, and most preferably 97%, or even 98% identity to a reference amino acid sequence (for example, the amino acid sequence shown in SEQ ID NOS: 1, 3, 5, 7 and 11)

or nucleic acid sequence (for example, the nucleic acid sequences shown in SEQ ID NOS:2, 4, 6, 8 and 12, respectively). For polypeptides, the length of comparison sequences will generally be at least 16 amino acids, preferably at least 20 amino acids, more preferably at least 25 amino acids, and most preferably 35 amino acids. For nucleic acids, the length of comparison sequences will generally be at least 50 nucleotides, preferably at least 60 nucleotides, more preferably at least 75 nucleotides, and most preferably 110 nucleotides.

Sequence identity is typically measured using sequence analysis software (for example, Sequence Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, Wis. 53705, BLAST, or PILEUP/PRETTYBOX programs). Such software matches identical or similar sequences by assigning degrees of homology to various substitutions, deletions, and/or other modifications. Conservative substitutions typically include substitutions within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid, asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine.

By an "isolated polypeptide" is meant a P450 polypeptide (for example, a CYP71D20 (SEQ ID NO:1), CYP71D21 (SEQ ID NO:3), CYP73A27 (SEQ ID NO:5), CYP73A28 (SEQ ID NO:7), or CYP92A5 (SEQ ID NO:11) polypeptide) that has been separated from components that naturally accompany it. Typically, the polypeptide is isolated when it is at least 60%, by weight, free from the proteins and naturally-occurring organic molecules with which it is naturally associated. Preferably, the preparation is at least 75%, more preferably at least 90%, and most preferably at least 99%, by weight, a P450 polypeptide. An isolated P450 polypeptide may be obtained, for example, by extraction from a natural source (for example, a plant cell); by expression of a recombinant nucleic acid encoding a P450 polypeptide; or by chemically synthesizing the protein. Purity can be measured by any appropriate method, for example, column chromatography, polyacrylamide gel electrophoresis, or by HPLC analysis.

By "derived from" or "obtained from" is meant isolated from or having the sequence of a naturally-occurring sequence (e.g., cDNA, genomic DNA, synthetic, or combination thereof).

By "isolated nucleic acid molecule" is meant a nucleic acid molecule, e.g., a DNA molecule, that is free of the nucleic acid sequence(s) which, in the naturally-occurring genome of the organism from which the nucleic acid molecule of the invention is derived, flank the nucleic acid molecule. The term therefore includes, for example, a recombinant DNA that is incorporated into a vector; into an autonomously replicating plasmid or virus; or into the genomic DNA of a prokaryote or eukaryote; or that exists as a separate molecule (for example, a cDNA or a genomic or cDNA fragment produced by PCR or restriction endonuclease digestion) independent of other sequences. The term "isolated nucleic acid molecule" also includes a recombinant DNA which is part of a hybrid gene encoding additional polypeptide sequences.

By "specifically hybridizes" is meant that a nucleic acid sequence is capable of hybridizing to a DNA sequence at least under low stringency conditions, and preferably under high stringency conditions. For example, high stringency conditions may include hybridization at approximately 42° C. in about 50% formamide, 0.1 mg/mL sheared salmon sperm DNA, 1% SDS, 2×SSC, 10% Dextran sulfate, a first wash at approximately 65° C. in about 2×SSC, 1% SDS, followed by a second wash at approximately 65° C. in about 0.1×SSC.

Alternatively high stringency conditions may include hybridization at approximately 42° C. in about 50% formamide, 0.1 mg/mL sheared salmon sperm DNA, 0.5% SDS, 5×SSPE, 1× Denhardt's, followed by two washes at room temperature in 2×SSC, 0.1% SDS, and two washes at between 55-60° C. in 0.2×SSC, 0.1% SDS. Reducing the stringency of the hybridization conditions may involve lowering the wash temperature and/or washing at a higher concentration of salt. For example, low stringency conditions may include washing in 2×SSC, 0.1% SDS at 40° C.

By "transformed cell" is meant a cell into which (or into an ancestor of which) has been introduced, by means of recombinant DNA techniques, a DNA molecule encoding (as used herein) a P450 polypeptide.

By "positioned for expression" is meant that the DNA molecule is positioned adjacent to a DNA sequence which directs transcription and translation of the sequence (i.e., facilitates the production of, for example, a P450 polypeptide, a recombinant protein, or an RNA molecule).

By "reporter gene" is meant a gene whose expression may be assayed; such genes include, without limitation, beta-glucuronidase (GUS), luciferase, chloramphenicol transacetylase (CAT), green fluorescent protein (GFP), beta-galactosidase, herbicide resistant genes, and antibiotic resistance genes.

By "expression control region" is meant any minimal sequence sufficient to direct transcription. Included in the invention are promoter elements that are sufficient to render promoter-dependent gene expression controllable for cell-, tissue-, or organ-specific gene expression, or elements that are inducible by external signals or agents (for example, light-, pathogen-, wound-, stress-, or hormone-inducible elements or chemical inducers such as salicylic acid (SA) or 2,2-dichloro isonicotinic acid (INA)); such elements may be located in the 5' or 3' regions of the native gene or engineered into a transgene construct.

By "operably linked" is meant that a gene and a regulatory sequence(s) are connected in such a way as to permit gene expression when the appropriate molecules (for example, transcriptional activator proteins) are bound to the regulatory sequence(s).

By "plant cell" is meant any self-propagating cell bounded by a semi-permeable membrane and typically is one containing a plastid. Such a cell also requires a cell wall if further propagation is desired. Plant cell, as used herein includes, without limitation, algae, cyanobacteria, seeds, suspension cultures, embryos, meristematic regions, callus tissue, leaves, roots, shoots, gametophytes, sporophytes, pollen, and microspores.

By "plant component" is meant a part, segment, or organ obtained from an intact plant or plant cell. Exemplary plant components include, without limitation, somatic embryos, leaves, stems, roots, flowers, tendrils, fruits, scions, and rootstocks.

By "transgene" is meant any piece of DNA which is inserted by artifice into a cell and typically becomes part of the genome, for example, the nuclear or plastidic genome, of the organism which develops from that cell. Such a transgene may include a gene which is partly or entirely heterologous (i.e., foreign) to the transgenic organism, or may represent a gene homologous to an endogenous gene of the organism.

By "transgenic" is meant any cell which includes a DNA sequence which is inserted by artifice into a cell and becomes part of the genome of the organism which develops from that cell. As used herein, the transgenic organisms are generally transgenic plants and the DNA (transgene) is inserted by

artifice into the nuclear or plastidic genome. A transgenic plant according to the invention may contain one or more engineered traits.

By "pathogen" is meant an organism whose infection of viable plant tissue elicits a disease response in the plant tissue. Such pathogens include, without limitation, bacteria, mycoplasmas, fungi, insects, nematodes, viruses, and viroids. Plant diseases caused by these pathogens are described in Chapters 11-16 of Agrios, Plant Pathology, 3rd ed., Academic Press, Inc., New York, 1988.

By "increased level of resistance" is meant a greater level of resistance to a disease-causing pathogen in a transgenic plant (or cell or seed thereof) of the invention than the level of resistance relative to a control plant (for example, a non-transgenic plant). In preferred embodiments, the level of resistance in a transgenic plant of the invention is at least 20% (and preferably 30% or 40%) greater than the resistance of a control plant. In other preferred embodiments, the level of resistance to a disease-causing pathogen is 50% greater, 60% greater, and more preferably even 75% or 90% greater than a control plant; with up to 100% above the level of resistance as compared to a control plant being most preferred. The level of resistance is measured using conventional methods. For example, the level of resistance to a pathogen may be determined by comparing physical features and characteristics (for example, plant height and weight, or by comparing disease symptoms, for example, delayed lesion development, reduced lesion size, leaf wilting and curling, water-soaked spots, and discoloration of cells) of transgenic plants.

By "purified antibody" is meant antibody which is at least 60%, by weight, free from proteins and naturally-occurring organic molecules with which it is naturally associated. Preferably, the preparation is at least 75%, more preferably 90%, and most preferably at least 99%, by weight, antibody, for example, an acquired resistance polypeptide-specific antibody. A purified P450 antibody may be obtained, for example, by affinity chromatography using a recombinantly-produced P450 polypeptide and standard techniques.

By "specifically binds" is meant an antibody which recognizes and binds a P450 protein but which does not substantially recognize and bind other molecules in a sample, for example, a biological sample, which naturally includes a P450 protein such as CYP71D20, CYP71D21, CYP73A27, CYP73A28, or CYP92A5.

Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof, and from the claims.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a schematic diagram of a proposed alternative pathway for the biosynthesis of capsidiol in elicitor-treated *Nicotiana tabacum* cells. 5-epi-aristolochene is synthesized from FPP by the action of a sesquiterpene cyclase, 5-epi-aristolochene synthase (EAS), and is subsequently hydroxylated at C1 and C3 to form capsidiol.

FIG. 2 is a graph showing an induction time course for sesquiterpene cyclase enzyme activity and sesquiterpene hydroxylase activity in cellulase-treated cell cultures. Sesquiterpene cyclase (5-epi-aristolochene synthase, EAS) enzyme activity was determined in extracts prepared from control (open squares) and elicitor-treated (closed squares) cells collected at the indicated time points. Sesquiterpene hydroxylase activity was determined using an indirect assay for control (open circles) and elicitor-treated (closed circles) cells. Cell cultures were incubated with [³H]-5-epi-aristolochene for 3 hours ending at the indicated time points

before quantifying the incorporation of radioactivity into extracellular capsidiol, a dihydroxylated form of aristolochene (Mandujano-Chávez et al., Arch. Biochem. Biophys. 381:285-294, 2000).

FIGS. 3A-3B are a series of graphs showing the dose dependent inhibition of 5-epi-aristolochene hydroxylase activity, by ancyimidol and ketoconazole. Cell cultures were incubated in the presence of cellulase (0.5 µg/mL) plus the indicated concentrations of ancyimidol (A) or ketoconazole (B) for 12 hours prior to measuring the in vivo 5-epi-aristolochene hydroxylase activity in the cell suspension cultures (squares), or the EAS enzyme activity in extracts prepared from the collected cells (triangles). The in vitro activity of a purified EAS preparation (Back and Chappell, J. Biol. Chem. 270:7375-7381, 1995) was also measured at the indicated inhibitor concentrations as an additional test for non-specific effects of these inhibitors (circles).

FIG. 4A is a schematic diagram of the primary structure of a generalized cytochrome P450 with conserved domains used for the design of PCR primers highlighted (SEQ ID NOS:26-29).

FIG. 4B is a list of the degenerate P450-specific primers (SEQ ID NOS:30-36) that were used in various combinations with vector specific primers in the amplification of cytochrome P450 cDNA fragments.

FIG. 4C is a scanned image of an ethidium bromide-stained agarose gel showing the PCR products amplified from a directional cDNA library prepared with mRNA isolated from elicitor-treated cells using the degenerate primer GRRXCP (A/G)- for (SEQ ID NO:35) and the T7 vector-specific primer (SEQ ID NO:37). The T3 vector-specific primer is also shown (SEQ ID NO:38).

FIG. 5 is a series of Northern blots showing the induction time course for CYP71D, CYP73A, CYP82E, CYP92A, and EAS transcript accumulation in elicitor treated cells. Total RNA was extracted from tobacco suspension cells incubated with the cellulase elicitor for the indicated durations, size fractionated by agarose gel electrophoresis under denaturing conditions, and transferred to a nylon membrane before probing with the respective full-length cDNAs. The uniformity of sample loading was verified by ethidium bromide staining of ribosomal RNA (Loading control).

FIGS. 6A-6B are a series of graphs showing carbon monoxide (CO) difference spectra of the microsomal fraction isolated from yeast expressing the CYP92A5 (A) and CYP71D20 (B) cDNAs. Expression of the respective plasmid constructs engineered into the yeast (WAT11) cells was induced by a galactose treatment, followed by isolation of microsomal preparations. The difference adsorption spectra of microsomes incubated in the presence (solid lines) and absence (broken lines) of carbon monoxide was determined.

FIGS. 7A-7D are a series of gas chromatograms of the reaction products formed upon incubation of microsomes isolated from WAT11 yeast cells containing the CYP71D20 expression construct (A and C) or vector control DNA (B and D) with sesquiterpene substrates. Microsomes isolated from the indicated yeast lines were incubated with 5-epi-aristolochene (A and B) or 1-deoxycapsidiol (C and D) in the presence (solid lines) or absence (dashed lines) of NADPH. The identities of 5-epi-aristolochene, 1-deoxycapsidiol, and capsidiol were verified by mass spectrometry.

FIGS. 8A-8D provide a sequence comparison of the amino acid sequence of *Nicotiana tabacum* 5-epi-aristolochene (sesquiterpene) hydroxylase NtCYP71D20 (SEQ ID NO:1) with other plant terpene hydroxylases (SEQ ID NOS:39-43). NtCYP71A5v1 (GenBank accession number CAA70575) catalyzes the mono-hydroxylation of nerol and geraniol, lin-

ear monoterpenes, while PaCYP71A1 (A35867) catalyzes the epoxidation of these substrates (Hallahan et al., Biochim. Biophys. Acta. 1201:94-100, 1994). MsCYP71D18 (AAD44150) and MpCYP71D13 (AAD44151) catalyze the mono-hydroxylation at C6 and C3 of limonene, a cyclic monoterpene, respectively (Lupien et al., Arch. Biochem. Biophys. 368:181-192, 1999). AtCYP701A3 (AAC39505) encodes for kaurene oxidase, which catalyzes a 3-step reaction including a hydroxylation followed by oxidation of a diterpene (Helliwell et al., Plant Physiol. 119:507-510, 1999). Shown are sequences from *Mentha piperita* (MpCYP71D13; SEQ ID NO:39), *Mentha spicata* (MsCYP71D18; SEQ ID NO:40), *Nepeta racemosa* (NrCYP71A5v1; SEQ ID NO:41), *Nicotiana tabacum* (NtCYP71D20; SEQ ID NO:1), *Persea americana* (PaCYP71A1; SEQ ID NO:42), and *Arabidopsis thaliana* (CYP701A3; SEQ ID NO:43). Conserved residues are shaded.

DETAILED DESCRIPTION

Capsidiol is a bicyclic, dihydroxylated sesquiterpene produced by several Solanaceous species in response to a variety of environmental stimuli. It is the primary antimicrobial compound produced by *Nicotiana tabacum* in response to fungal elicitation, and it is formed via the isoprenoid pathway from 5-epi-aristolochene. Much of the biosynthetic pathway for the formation of this compound has been elucidated, except for the enzyme(s) responsible for the conversion of the allylic sesquiterpene 5-epi-aristolochene to its dihydroxylated form, capsidiol.

Accordingly, an in vivo assay for 5-epi-aristolochene hydroxylase-activity was developed and used to demonstrate a dose dependent inhibition of activity by ancyimidol and ketoconazole, two well-characterized inhibitors of cytochrome P450 enzymes. Using degenerate oligonucleotide, primers designed to the well-conserved domains found within most P450 enzymes, including the heme binding domain, cDNA fragments representing four distinct P450 families (CYP71, CYP73, CYP82, and CYP92) were amplified from a cDNA library prepared against mRNA from elicitor-treated cells using PCR. The PCR fragments were subsequently used to isolate full-length cDNAs (CYP71D20 (SEQ ID NO:2) and D21 (SEQ ID NO:4), CYP73A27 (SEQ ID NO:6) and A28 (SEQ ID NO:8), CYP82E1 (SEQ ID NO:10), and CYP92A5 (SEQ ID NO:12)), and these in turn were used to demonstrate that the corresponding mRNAs were all induced in elicitor-treated cells, albeit with different induction patterns.

EXAMPLES

There now follows a description of the cloning of several P450s from *Nicotiana tabacum*. These examples are provided for the purpose of illustrating the invention, and are not to be considered as limiting.

Inhibition of the 5-Epi-Aristolochene to Capsidiol Conversion by P450 Antagonists

Using an indirect assay, a detailed induction time course of 5EAH activity in elicitor-induced cell cultures was determined relative to that of EAS activity (FIG. 2), the well-characterized sesquiterpene cyclase activity that catalyzes the formation of 5-epi-aristolochene from FPP (FIG. 1). Using assays for EAS and 5EAH, EAS activity is not detectable in control cell cultures, but is induced significantly within 3 hours and reaches its maximal level within 15 to 18 hours of elicitor-treatment. Similar to the EAS enzyme activity, 5EAH

activity was negligible in control cell cultures. Nonetheless, after an apparent lag phase of 8 hours, a rapid induction of hydroxylase activity was observed 10 to 15 hours post elicitor addition to the cell cultures, reaching a maximum by 18 hours followed by a rather gradual decline of 10 to 20% over the next 8 hours.

Tobacco cell suspension cultures treated with cellulase plus varying concentrations of ancymidol or ketoconazole were pre-incubated for 12 hours before measuring the cells' ability to convert exogenous supplied [³H] labeled 5-epi-aristolochene to radiolabeled capsidiol during a subsequent 3 hour incubation period (FIGS. 3A-3B). Apparent activity of 5EAS was inhibited in a dose-dependent manner with approximately 50% inhibition by either 25 μ M ancymidol or ketoconazole, and more than 80% by 75 μ M ancymidol and 95% by 100 μ M ketoconazole (FIGS. 3A and 3B). Importantly, neither the in vitro activity of recombinant EAS nor the induction of EAS in the elicitor-treated cell cultures was significantly affected by ancymidol at concentrations as high as 100 μ M (FIG. 3A). Ketoconazole also does not appear to affect the in vitro activity of EAS. However, the inducibility of cyclase activity in elicitor-treated cell extracts was inhibited by ketoconazole at concentrations above 50 μ M (FIG. 3B). Therefore, the specificity of ketoconazole as an inhibitor of P450 type reactions should be assessed at or below a concentration of 50 μ M under these experimental conditions. Isolation of Elicitor-Inducible Cytochrome P450 cDNAs

A two-step approach for the isolation of candidate P450 cDNAs was followed. A PCR strategy was first employed using a directional cDNA library prepared against mRNA isolated from elicitor-induced cells as the template and degenerate PCR primers (FIGS. 4A-4C). Sequence alignments of cytochrome P450s from multiple families across kingdoms were used to identify conserved regions to which a series of degenerate primers were prepared (FIGS. 4A and 4B). In cloning experiments, 450 to 550 bp products were expected from reactions utilizing the primer prepared to the heme-binding domain (GRRXCP(A/G)) (SEQ ID NOS:27 and 28) and the T7 vector primer (FIG. 4C). The mixtures of reaction products were shotgun cloned, and approximately 100 of the cloned PCR fragments were sequenced. About half of the sequenced DNAs contained signature sequences typical of P450 enzymes as revealed by BlastX database searches, and these corresponded to typical plant P450 family members of the CYP71, CYP73, CYP92 and CYP82 classes. Each of these PCR fragments was isolated multiple times in separate experiments. In addition, we isolated full-length cDNAs for these P450 family members. Table 1 compares the similarity and identity of the full-length cDNAs of P450 family members with those of their nearest family member in the GenBank database. In addition, FIGS. 8A-8D shows an amino acid alignment of several terpene cytochrome P450s. Alignments were performed using the algorithm of the MACVECTOR software suite.

TABLE 1

Full-length cDNAs cloned from an elicited cDNA library			
Cytochrome P450 cDNA clone	Nearest relative/ accession number	% Identity	% Similarity
CYP71D20	CYP71D7 (<i>S. chacoense</i>) Gen EMBL U48435	76.5	88.8
CYP71D21	CYP71D7 (<i>S. chacoense</i>) Gen EMBL U48435	76.3	88.8
CYP73A27	CYP73A15 (<i>P. vulgaris</i>) Gen EMBL Y09447	79.4	92.6

TABLE 1-continued

Full-length cDNAs cloned from an elicited cDNA library			
Cytochrome P450 cDNA clone	Nearest relative/ accession number	% Identity	% Similarity
CYP73A28	CYP73A15 (<i>P. vulgaris</i>) Gen EMBL Y09447	79.2	92.4
CYP82E1	CYP82E1 (<i>N. tabacum</i>) Gen EMBL AB015762	100.0	100.0
CYP92A5	CYP92A3 (<i>N. tabacum</i>) Gen EMBL X96784	95.5	98.6

The cloned fragments were used in a second step to isolate full-length clones from the cDNA library. Screening the cDNA library by hybridization with the CYP71 and CYP73 gene fragments yielded four full-length cDNAs, two CYP71Ds and two CYP73As. The former clones were designated CYP71D20 and CYP71D21, and the latter were designated CYP73A27 and CYP73A28. The other two cDNA fragments corresponded to tobacco cDNAs already found in the GenBank database, CYP82E1 and CYP92A3. These two cDNAs were cloned using specific primers designed with the help of the available sequence information to amplify the full-length cDNA.

Induction of Cytochrome P450 mRNAs in Elicitor-Treated Cells

To correlate a biochemical role for P450s in sesquiterpene metabolism, RNA blot analyses were used to determine the steady-state levels of the mRNAs coding for all four of the cytochrome P450 clones and EAS in control and elicitor-treated cells (FIG. 5). The mRNAs for all four of the P450s were rapidly and transiently induced with slightly different time courses relative to one another and to the EAS mRNA. CYP73A27 mRNA, for instance, displayed an induction pattern similar to that of EAS with the maximum mRNA level occurring 9 to 12 hours after elicitation. While the EAS mRNA remained high throughout the duration of the experiment, the CYP73A27 mRNA was negligible in cells 24 hours after elicitor-treatment. In contrast, the CYP71D mRNA was more rapidly induced than the EAS mRNA, reached its maximum 6 to 9 hours after elicitation, and was declining by 12 hours when the EAS mRNA level was still very high.

Functional Identification of CYP71D20 as 5-Epi-Aristolochene Hydroxylase

To ascribe functional identity to the various P450 cDNAs, full-length cDNAs for CYP71D20, CYP82E1 and CYP92A5 were inserted into the yeast expression vector pYedP60 (Urban et al., *Biochimie* 72:463-472, 1990; Pompon et al., *Methods Enzymol.* 272:51-64, 1996) and the expression of each in WAT11, a yeast line containing an integrated *Arabidopsis thaliana* cytochrome reductase gene (Pompon et al., *Methods Enzymol.* 272:51-64, 1996; Urban et al., *J. Biol. Chem.* 272:19176-19186, 1997), was determined. Engineering the CYP73A27 cDNA required an extra modification because of an unusually long N-terminus with several hydrophilic residues that may interfere with proper intracellular targeting (Nedelkina et al., *Plant Mol. Biol.* 39:1079-1090, 1999). This unusual leader sequence therefore was replaced with the membrane anchoring sequence of CYP73A1, a cinnamate 4-hydroxylase previously demonstrated to express well in yeast (Fahrendorf and Dixon, *Arch. Biochem. Biophys.* 305:509-515, 1993; Pompon et al., *Methods Enzymol.* 272:51-64, 1996). Expression of all these cDNAs was under the control of the glucose-repressible, galactose-inducible GAL10-CYC1 promoter (Guarente et al., *Proc. Natl. Acad. Sci.*

U.S.A. 79:7410-7414, 1982), and expression was compared to yeast transformed with the parent pYeDP60 vector (control) alone.

After induction with galactose for approximately 16 hours, control cells and cells containing the various P450 constructs were collected, and microsomes prepared from each were analyzed for general P450 expression by CO-difference spectroscopy (Omura and Sato, J. Biol. Chem. 239:2370-2378, 1964). Microsomes prepared from cells containing the CYP71D20 (FIG. 6A) and CYP92A5 (FIG. 6B) constructs both showed characteristic CO difference spectra with peaks at 450 nm, indicating that the encoded proteins were assembling properly with their heme cofactor. Using the extinction coefficient of $91 \text{ mM}^{-1}\cdot\text{cm}^{-1}$ for heme binding proteins (Omura and Sato, J. Biol. Chem. 239:2370-2378, 1964), it was determined that approximately 107 pmol of CYP71D20 and 268 pmol of CYP92A5 were expressed in the yeast cells per milligram of total yeast protein.

Both 5-epi-aristolochene and 1-deoxycapsidiol were metabolized to only one product with the same retention time as capsidiol. Obvious by its absence, no reaction product having a retention time similar to deoxycapsidiol was detectable in the 5-epi-aristolochene incubations (FIGS. 7A-7D). Co-injection of authentic capsidiol with the respective reaction products resulted in a single GC peak having a 16.2 minute retention time, identical to capsidiol. Mass spectra patterns for the separate reaction products were identical to that for the capsidiol standard (EIMS m/z 236, 221, 203, 185, 175, 163, 157, 133, 121, 107, 93, 79, 67, 55, 43, 41).

The in vivo assay data presented in FIGS. 2 and 3A-3B of the current work indicate that the conversion of 5-epi-aristolochene is catalyzed by at least one inducible cytochrome P450 mediated reaction.

Furthermore, any of the cytochrome p450 polypeptides described herein may include one or more hydroxylase activities which can incorporate hydroxyl groups into at least two distant sites on an isoprenoid compound. The addition of these hydroxyl groups may occur, for example, sequentially, by adding a hydroxyl group first to one site and then the other, in either order. Moreover, such hydroxylases may be mutated to limit their ability to hydroxylate a substrate at only one site, or, alternatively, to provide stereochemical specificity to their hydroxylating activity.

The above-described experiments were performed using the following materials and methods.

Chemicals

Standard laboratory reagents were purchased from Becton Dickinson Microbiology Systems (Sparks, Md.), FisherBiotech (Fair Lawn, N.J.) and Sigma Chemical Company (St. Louis, Mo.).

Biological Materials and Induction Treatments

Nicotiana tabacum cv. KY14 plants and cell suspension cultures were used. Cell suspension cultures were maintained in modified Murashige-Skoog (Vögeli and Chappell, Plant Physiol. 88:1291-1296, 1988). Cultures in their rapid phase of growth (3 days old) were used for all experiments. At the indicated times, cells were collected and separated from media by vacuum filtration and stored at -80°C .

Induction treatments were performed by the addition of the fungal elicitors, cellulase (*Trichoderma viride*, Type RS, Onozuka) or paraciticein (O'Donohue et al., Plant Mol. Biol. 27:577-586, 1995) at the indicated concentrations. Paraciticein was purified from *E. coli* cells overexpressing a recombinant paraciticein protein containing a carboxy-terminal histidine purification tag.

In vivo 5-Epi-Aristolochene Hydroxylase Assay and Inhibition Studies

5-epi-aristolochene hydroxylase-activity was measured as the incorporation of [^3H]-5-epi-aristolochene into extracellular capsidiol by intact cells. [^3H]-5-epi-aristolochene was produced by incubating an excess of [^3H] farnesyl diphosphate ($1 \mu\text{M}$, 20.5 Ci/mmol) with recombinant 5-epi-aristolochene synthase (Back et al., Arch. Biochem. Biophys. 315:527-532, 1994; Rising et al., J. Am. Chem. Soc. 122: 1861-1866, 2000). The hexane extractable radioactivity from reactions was treated with a small amount of silica to remove any farnesol or residual FPP before quantifying the yield of radioactive 5-epi-aristolochene by liquid scintillation counting. The hexane solvent was removed under a gentle stream of N_2 gas, and the dried residue was re-dissolved in acetone. Control and elicitor-treated cells were then incubated with [^3H]-5-epi-aristolochene (approximately 100,000 dpm at 2.5 nM) for 3 hour periods at various points during an induction time course before collecting the cell and media samples. Detection and quantification of capsidiol in the extracellular culture media was performed as reported previously (Chappell et al., Phytochemistry 26:2259-2260, 1987), and the amount of radioactivity incorporated into capsidiol was determined. For these determinations, samples were separated by TLC, and the zones corresponding to capsidiol were scraped from the plate for scintillation counting.

Inhibition studies were performed by the addition of the P450 inhibitors ancymidol (Coolbaugh et al., Plant Physiol. 62:571-576, 1978; Hoshino et al., Phytochemistry 38:609-613, 1995) and ketoconazole (Hoshino et al., Phytochemistry 38:609-613, 1995; Rademacher, Annu. Rev. Plant Physiol. Plant Mol. Biol. 51:501-531, 2000) directly to the cell cultures or enzyme assay mix. Cell cultures were incubated in the presence of cellulase ($0.5 \mu\text{g/mL}$) and indicated concentrations of ancymidol or ketoconazole for 12 hours prior to the addition of [^3H]-5-epi-aristolochene. After a further 3 hour incubation period, the cells and media were collected. The amount of radioactivity incorporated into extracellular capsidiol was determined as described above. To evaluate secondary effects of these inhibitors, the level of inducible sesquiterpene cyclase activity in the collected cells was determined according to Vögeli et al. (Plant Physiol. 93:182-187, 1990), as well as in vitro assays with purified recombinant EAS (Back et al., Arch. Biochem. Biophys. 315:527-532, 1994) incubated with the indicated concentrations of ancymidol and ketoconazole.

All experiments were replicated in several independent trials. While the absolute values presented may have varied between experiments by as much as 50%, the trends and time courses were consistent throughout.

Construction of an Elicitor-Induced cDNA Library

Cell cultures were incubated with fungal elicitor ($0.5 \mu\text{g}$ cellulase/mL) for 6 hours before collecting the cells by filtration. The cells were kept frozen at -80°C until total RNA was extracted from them using Trizol (Life Technologies, Rockville, Md.) according to the manufacturer's instructions. Poly (A) $^+$ RNA was purified by two rounds of oligo (dT) cellulose column chromatography (Life Technologies, Rockville, Md.). cDNA synthesis and library construction were subsequently carried out using the UNI-ZAP XR library kit (Stratagene, La Jolla, Calif.), according to manufacturer's instructions.

PCR Cloning Strategy

Cytochrome P450 cDNA fragments were amplified from the elicitor-induced cDNA library using various combinations of degenerate forward and reverse primers with the vector-specific T3 and T7 primers. The template DNA was

13

prepared from a 500 μ L aliquot of the elicitor-induced cDNA library (3×10^6 pfu/ μ L) by heat denaturation at 70° C. for 10 minutes, followed by phenol/chloroform extraction, ethanol precipitation and re-suspension in 500 μ L of sterile, deionized water. Amplification reactions were performed in 50 μ L volumes containing 50 mM KCl; 10 mM Tris-HCl, pH 8.8; 1.5 mM $MgCl_2$; 200 μ M of each dNTP; 2 μ L template DNA; 20 pmol each of forward and reverse primer; and 1 unit Taq Polymerase (Life Technologies, Rockville, Md.). Reactions were preheated at 94° C. for 2 minutes, followed by thirty-five cycles of denaturing at 94° C. for 1 minute, annealing at 50° C. for 1 minute 30 seconds, and polymerization at 72° C. for 2 minutes. The reactions were completed by a 10-minute extension at 72° C. Aliquots of the reaction products were examined for DNA products by agarose gel fractionation, and ligated directly into the pGEM-T Easy vector (Promega, Madison, Wis.). Resulting recombinant plasmids containing insert DNAs within the expected size range were sequenced using T7 and Sp6 primers.

DNA Sequencing

All the DNA sequencing reactions were performed using the BIGDYE™ Terminator Cycle sequencing kit (Perkin-Elmer, Wellesley, Mass.) with the sequences being read on an automated ABI Prism 310 Genetic Analyzer (Applied Biosystems, Foster City, Calif.). Computer assessment of the DNA sequence information was performed using the MACVECTOR (Oxford Molecular, Madison, Wis.) software package.

cDNA Library Screening

The cDNA library was screened with digoxigenin labeled probes. A 258 bp DNA fragment amplified from the pGEM-deg6.4 clone using gene-specific forward (5'-GGCG-GAGAATTGTCCTGGAATGTCATTGGTTTGG-3' (SEQ ID NO:13)) and reverse (5'-GTACAATAGTGAGGT-TGACAATG-3' (SEQ ID NO:14)) primers; and a 374 by DNA fragment amplified from the pBKS-CYPB3.843 clone with specific forward (5'-GGTGGTTGTGAATGCATG-3' (SEQ ID NO:15)) and reverse (5'-TTATGCAGCAATAG-GCTTGAAGACA-3' (SEQ ID NO:16)) primers, were used to screen for CYP71Ds. The probes were labeled with digoxigenin-11-dUTP using the PCR DIG Labeling Mix (Roche Molecular Biochemicals, Indianapolis, Ind.), hybridized to plaque lifts of the cDNA library plated at approximately 10,000 PFUs per 150 mm plate, and was hybridization detected with the DIG detection system according to the manufacturer's instructions (Roche Molecular Biochemicals, Indianapolis, Ind.). Plaques exhibiting strong hybridization were plaque purified, auto subcloned to their plasmid forms according to the manufacturer's recommendations (Stratagene, La Jolla, Calif.), and then subjected to DNA sequencing as described above.

RNA Analysis

RNA gel blot analysis was carried out using 10 vtg aliquots of total RNA. RNA samples were heat-denatured at 70° C. for 15 minutes in sample buffer (1 \times MOPS, 50% formamide, 16% formaldehyde, 30% glycerol, and 3% ethidium bromide), and size fractionated on a 1.2% agarose gel containing 1 \times MOPS and 18.1% formaldehyde. Uniformity of sample loading was determined by visual inspection of the gel for rRNA bands. The RNAs were then transferred to a Zeta Probe nylon membrane (Bio-Rad Laboratories, Hercules, Calif.) and hybridized according to the manufacturer's recommendations. Full-length cDNA probes were labeled with [32 P]-dCTP (PRIME-IT Kit, Stratagene, La Jolla, Calif.) prior to hybridization. After hybridization, the membranes were washed in 2 \times SSC/0.1% SDS once at room temperature followed by sequential washes in 0.2 \times SSC/0.1% SDS at 42° C.

14

and 65° C. Hybridization was detected with a Phosphorimager (Molecular Dynamics, model 445 SI).

Construction of Yeast Expression Vectors

The coding regions of the P450 cDNAs were cloned into the pYeDP60 expression vector (Urban et al., J. Biol. Chem. 272:19176-19186, 1990; Pompon et al., Methods Enzymol. 272:51-64, 1996). Appropriate BamHI, EcoRI, and SstI restriction sites (underlined) were introduced via PCR primers containing these sequences either upstream of the translation start site (ATG) or downstream of the stop codon (TAA or TGA). The primers used to amplify the CYP71D20 cDNA were 5'-GGG GGATCCATGCAATTCTTCAGCTTGGTTTCC-3' (SEQ ID NO:17) and 5'-GGG GAATTCCTTACTCTCGAGAAGGTTGATAAGG-3' (SEQ ID NO:18); for the CYP82E1 cDNA 5'-CCC GGATCCATGTATCATCTTCTTTCTCCC-3' (SEQ ID NO:19) and 5'-GGG GAATTCCTCAATATTGATAAAGCGTAGGAGG-3' (SEQ ID NO:20); and for the CYP92A3 cDNA 5'-CCC GGATCCATGCAATCCTTCAGCTTGGTTTCC-3' (SEQ ID NO:21) and 5'-GGG GAGCTCTCACTCGCAAGAAGATTGATAAGG-3' (SEQ ID NO:22). Two long, overlapping (italicized) primers 5'-GCCATTATCGGCGCAATACTAATCTC-CAAACTCCGCGGTAAAAAATTC AAGC TCCCAC-CTGGTCCAACAGCAGTC-3' (SEQ ID NO:23) and 5'-GGG GGATCCATGGACCTCCTCCTCATAGAAAAA-CCCTCGTCGCCTTATTC GCCGCCATTATCGGCG-CAATACTA-3' (SEQ ID NO:24) coding for the N-terminal sequence of CYP73A1 (GenEMBL Z17369) up to the hinge region were used for the modification of the membrane anchoring segment of CYP73A27 to avoid possible problems with intracellular targeting due to the unusual N-terminus (Nedelkina et al., 1999); the reverse primer used for both amplifications was 5'-GGG GAGCTCTTATGCAGCAATAGGCTTGAAGAC-3' (SEQ ID NO:25). CYP71D20 and CYP73A27 were amplified using full-length cDNA templates, whereas CYP82E1 and CYP92A5 were amplified directly from the cDNA library template. Amplifications were performed in 50 μ L reactions containing 1 \times Pfx amplification buffer; 1 mM $MgSO_4$; 300 μ M of each dNTP; 10 ng template DNA; 20 pmol each of forward and reverse primer; and 1.25 units PLATINUM® Pfx Polymerase (Life Technologies, Rockville, Md.). Reactions were preheated at 94° C. for 2 minutes, followed by thirty-five cycles of denaturing at 94° C. for 15 seconds, annealing at 55° C. for 30 seconds, and elongating at 68° C. for 1.5 minutes. PCR products were ligated into the pGEM-T EASY vector (Promega, Madison, Wis.) and subcloned into the pYeDP60 vector. The resulting constructs were validated by a combination of PCR and DNA sequencing.

Yeast Expression Studies

Verified pYeDP60-P450 cDNA constructs were introduced into the yeast WAT11 line, a derivative of the W303-1B strain (MATa; ade 2-1; his 3-11; leu 2-3, -112; ura 3-1; can^R; cyr⁺), provided by Dr. P. Urban (Centre de Génétique Moléculaire, CNRS, Gif-sur-Yvette, France). The endogenous NADPH-cytochrome P450 reductase (CPR1) locus has been replaced with ATR1, a NADPH-cytochrome P450 reductase from *Arabidopsis thaliana* (Pompon et al., Methods Enzymol. 272:51-64, 1996; Urban et al., J. Biol. Chem. 272: 19176-19186, 1997), in the WAT11 line. Yeast was grown overnight in a 30° C. shaker in YPAD (1 g/l yeast extract; 1 g/L peptone; 20 g/L glucose; 200 mg/L adenine) liquid media. Cultures were harvested at an A_{600} between 0.5 and 1.5. Cells were collected by centrifugation at 2,500 \times g for 5 minutes at 4° C., and resus-

pended in ice-cold, sterile dH₂O. Cells were pelleted again as above and resuspended in 1M sorbitol. Forty μ L of yeast suspension was mixed with 0.5 to 1 μ g plasmid DNA (in <5 μ L dH₂O) in a pre-chilled 0.5 mL tube, and transferred to a chilled cuvette with a 0.2 cm electrode gap. One pulse at 1.5 kV, 25 μ F, and 200 Ohms was applied by an Eppendorf Electroporator (model 2510). A mixture of 500 μ L of YPAD/1M sorbitol was immediately added to the electroporated cells. Cells were allowed to recover at 30° C. for 1 hour, then spread onto SGI plates (1 g/l bactocasamino acids; 7 g/l yeast nitrogen base; 20 g/l glucose; 20 mg/l tryptophan; and 20 g/l agar). Transformed colonies appeared after 3 to 6 days of incubation at 30° C. Recombinant plasmids were confirmed by PCR assays performed directly on randomly selected yeast colonies.

For expression studies, one colony was added to SGI media (1 g/l bactocasamino acids; 7 g/l yeast nitrogen base; 20 g/l glucose; and 20 mg/l tryptophan) and grown at 30° C. for approximately 24 hours. An aliquot of this culture was diluted 1:50 into 250 mL of YPGE (10 g/l bactopectone; 10 g/l yeast extract; 5 g/l glucose; and 3% ethanol by volume) and the cells were grown until all glucose was consumed. The absence of glucose was determined by placing a 200 μ L aliquot of culture into a 1.5 mL tube, inserting a DIASTIX urinalysis reagent strip (Bayer, Elkhart, Ind.) for 30 seconds, and observing colorimetric changes indicating glucose levels. Induction was initiated by the addition of 5 grams of galactose (final concentration of 2%). The cultures were maintained at 30° C. for an additional 16 hours before collecting the cells by centrifugation at 7,000 \times g for 10 minutes. The pelleted cells were washed with 100 mL of TES buffer (50 mM Tris-HCl, pH 7.5; 1 mM EDTA; 0.6 M sorbitol). The cells were centrifuged as above, resuspended in 100 mL of TES-M (TES supplemented with 10 mM 2-mercaptoethanol), and allowed to incubate at room temperature for 10 minutes. The yeast cells were centrifuged again at 7,000 \times g for 10 minutes, and the pellet was resuspended in 2.5 mL extraction buffer (1% bovine serum albumin, fraction V; 2 mM 2-mercaptoethanol; 1 mM phenylmethylsulfonyl fluoride, all dissolved in TES). Glass beads (0.5 mm in diameter, Biospec Products, Inc., Bartlesville, Okla.) were added until skimming the surface of the cell suspension. Cell walls were disrupted manually by hand shaking in a cold room for 10 min at 30 second intervals separated by 30 second intervals on ice. Cell extracts were transferred to a 50 mL centrifuge tube, the glass beads were washed three times with 5 mL of extraction buffer, and the washes were pooled with the original cell extracts. Microsomes were prepared by differential centrifugation at 10,000 g for 10 minutes at 4° C. to remove cellular debris, followed by centrifugation at 100,000 \times g for 70 minutes at 4° C., and microsomal pellets were resuspended in 1.5 mL TEG-M buffer (50 mM Tris-HCl, pH 7.5; 1 mM EDTA; 20% glycerol; and 1.5 mM 2-mercaptoethanol) and stored frozen at -80° C. until further assayed.

CO Difference Spectra

Fe²⁺·CO vs. Fe²⁺ difference spectroscopy (Omura and Sato, J. Biol. Chem. 239:2370-2378, 1964) was performed using 0.4 mL of microsomes suspended in 1.6 mL of 50 mM Tris-HCl, pH 7.5; 1 mM EDTA; and 20% glycerol. A small amount of the reducing agent, sodium dithionite, was added, and the mixture was distributed between two cuvettes. A baseline was recorded between 400 and 500 nm on a Perkin Elmer Lambda 18 UV/visible spectrophotometer. CO was then bubbled into the sample cuvette for 1 minute, and the difference spectrum recorded again. The amount of functional P450 was estimated based on an absorbance coefficient of 91 mM⁻¹·cm⁻¹.

5-Epi-Aristolochene-1,3-Hydroxylase Assays

5-epi-aristolochene-1,3-hydroxylase assays were performed in 0.5 mL polyethylene tubes in 100 μ L volumes. 5-epi-aristolochene or 1-deoxycapsidiol dissolved in hexane was added to the tube, and the organic solvent was removed by incubation of the open tube at 30° C. 5-epi-aristolochene and 1-deoxycapsidiol were resuspended in 2 μ L dimethyl sulfoxide before adding the reaction mixture. Reactions were carried out in 100 mM Tris-HCl, pH 7.5, to which microsomal protein was added to a final concentration of 1 mg/mL. Reactions were initiated by the addition of 2 mM NADPH. The final concentration of 5-epi-aristolochene and 1-deoxycapsidiol in these assays varied from 20 to 50 μ M. After incubations for variable lengths of time at 30° C., the reactions were extracted with two volumes of ethyl acetate. The organic extracts were concentrated and evaluated by GC and GC-MS along with standards of 5-epi-aristolochene (Whitehead et al., *Phytochemistry* 28:775-779, 1989; Rising et al., *J. Am. Chem. Soc.* 122:1861-1866, 2000), 1-deoxycapsidiol (Whitehead et al., *Phytochemistry* 29:479-482, 1990), and capsidiol (Whitehead et al., *Phytochemistry* 26:1367-1369, 1987; Milat et al., *Phytochemistry* 30:2171-2173, 1991). GC analysis was routinely performed with an HP5890 GC equipped with a Hewlett-Packard HP-5 capillary column (30 m \times 0.25 mm, 0.25 μ m phase thickness) and FID as described previously (Rising et al., *J. Am. Chem. Soc.* 122:1861-1866, 2000). GC-MS analysis was performed at the University of Kentucky Mass Spectrometry Facility using a Varian 3400 gas chromatograph and a Finnigan INCOS 50 quadrupole mass selective detector. The GC was equipped with a J&W DB-5 ms capillary column (15 m \times 0.25 mm, 0.25 μ m phase thickness) and run with He as the carrier gas (10 psi.). Splitless injections were done at an injection port temperature of 280° C. The column temperature was maintained at 40° C. for 1 minute and then increased to 280° C. at 10° C. per minute. Following separation by the GC column, samples were introduced directly into the electron impact ionization source. Mass spectra were acquired at 70 eV, scanning from 40-440 Da in 1 second.

Production of Cytochrome P450s

Using the standard molecular techniques described herein, the isolation of additional cytochrome P450 coding sequences is readily accomplished. For example, using all or a portion of the amino acid sequence of any of the disclosed P450s, one may readily design P450-specific oligonucleotide probes, including P450 degenerate oligonucleotide probes (i.e., a mixture of all possible coding sequences for a given amino acid sequence). These oligonucleotides may be based upon the sequence of either DNA strand and any appropriate portion of the P450 nucleotide sequence. General methods for designing and preparing such probes are provided, for example, in Ausubel et al., 2000, *Current Protocols in Molecular Biology*, Wiley Interscience, New York, and Berger and Kimmel, *Guide to Molecular Cloning Techniques*, 1987, Academic Press, New York. These oligonucleotides are useful for P450 gene isolation, either through their use as probes capable of hybridizing to a P450 complementary sequence, or as primers for various amplification techniques, for example, polymerase chain reaction (PCR) cloning strategies.

Hybridization techniques and screening procedures are well known to those skilled in the art and are described, for example, in Ausubel et al. (supra); Berger and Kimmel (supra); Chen et al. (*Arch. Biochem. Biophys.* 324:255, 1995); and Sambrook et al. (*Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, New York). If desired, a combination of different oligonucleotide probes

may be used for the screening of a recombinant DNA library. The oligonucleotides may be detectably-labeled using methods known in the art and used to probe filter replicas from a recombinant DNA library. Recombinant DNA libraries are prepared according to methods well known in the art, for example, as described in Ausubel et al. (supra), or they may be obtained from commercial sources.

As discussed above, P450 oligonucleotides may also be used as primers in a polymerase chain reaction (PCR) amplification cloning strategy. PCR methods are well known in the art and are described, for example, in PCR Technology, Erlich, ed., Stockton Press, London, 1989; PCR Protocols: A Guide to Methods and Applications, Innis et al., eds., Academic Press, Inc., New York, 1990; and Ausubel et al. (supra). Primers are optionally designed to allow cloning of the amplified product into a suitable vector, for example, by including appropriate restriction sites at the 5' and 3' ends of the amplified fragment (as described herein). If desired, a P450 gene may be isolated using the PCR "RACE" technique, or Rapid Amplification of cDNA Ends (see, e.g., Innis et al. (supra)). By this method, oligonucleotide primers based on a P450 sequence are oriented in the 3' and 5' directions and are used to generate overlapping PCR fragments. These overlapping 3'- and 5'-end RACE products are combined to produce an intact full-length cDNA. This method is described in Innis et al. (supra) and Frohman et al. (Proc. Natl. Acad. Sci. U.S.A. 85:8998, 1988).

Additional methods for identifying sequences encoding P450s are provided in Maughan et al. (Arch. Biochem. Biophys. 341:104-111, 1997) and Clark et al. (Plant Mol. Biol. 33:875-885, 1997).

Useful P450 sequences may be isolated from any appropriate organism. Confirmation of a sequence's relatedness to a P450 polypeptide disclosed herein may be accomplished by a variety of conventional methods, for example, by comparing the sequence with a known P450 sequence found in a database. In addition, the activity of any P450 may be evaluated according to any of the techniques described herein.

P450 Polypeptide Expression

P450 polypeptides may be produced by transformation of a suitable host cell with all or part of a P450 DNA (for example, anyone of the P450 cDNAs described herein) in a suitable expression vehicle or with a plasmid construct engineered for increasing the expression of a P450 polypeptide in vivo.

Those skilled in the field of molecular biology will appreciate that any of a wide variety of expression systems may be used to provide the recombinant protein. The precise host cell used is not critical to the invention. The P450 protein may be produced in a prokaryotic host, for example, *E. coli* TB 1, or in a eukaryotic host, for example, *Saccharomyces cerevisiae*, insect cells, mammalian cells (for example, COS 1 or NIH 3T3 cells), or any of a number of plant cells including, without limitation, algae, tree species, ornamental species, temperate fruit species, tropical fruit species, vegetable species, legume species, monocots, dicots, or in any plant of commercial or agricultural significance. Particular examples of suitable plant hosts include, but are not limited to, Conifers, Petunia, Tomato, Potato, Tobacco, Grape, Arabidopsis, Lettuce, Sunflower, Oilseed rape, Flax, Cotton, Sugarbeet, Celery, Soybean, Alfalfa, Medicago, Lotus, Vigna, Cucumber, Carrot, Eggplant, Cauliflower, Horseradish, Morning Glory, Poplar, Walnut, Apple, Asparagus, Grape, Rice, Maize, Millet, Onion, Barley, Orchard grass, Oat, Rye, Tobacco and Wheat.

Such cells are available from a wide range of sources including: the American Type Culture Collection (Rockland,

Md.); or from any of a number of seed companies, for example, W. Atlee Burpee Seed Co. (Warminster, Pa.), Park Seed Co. (Greenwood, S.C.), Johnny Seed Co. (Albion, Me.), or Northrup King Seeds (Harstville, S.C.). Descriptions and sources of useful host cells are also found in Vasil I. K., Cell Culture and Somatic Cell Genetics of Plants, Vol I, II, III; Laboratory Procedures and Their Applications, Academic Press, New York, 1984; Dixon, R. A., Plant Cell Culture—A Practical Approach, IRL Press, Oxford University, 1985; Green et al., Plant Tissue and Cell Culture, Academic Press, New York, 1987; and Gasser and Fraley, Science 244:1293, 1989.

For prokaryotic expression, DNA encoding a P450 polypeptide is carried on a vector operably linked to control signals capable of effecting expression in the prokaryotic host. If desired, the coding sequence may contain, at its 5' end, a sequence encoding any of the known signal sequences capable of effecting secretion of the expressed protein into the periplasmic space of the host cell, thereby facilitating recovery of the protein and subsequent purification. Prokaryotes most frequently used are various strains of *E. coli*; however, other microbial strains may also be used. Plasmid vectors are used which contain replication origins, selectable markers, and control sequences derived from a species compatible with the microbial host. Examples of such vectors are found in Pouwels et al. (supra) or Ausubel et al. (supra). Commonly used prokaryotic control sequences (also referred to as "regulatory elements") are defined herein to include promoters for transcription initiation, optionally with an operator, along with ribosome binding site sequences. Promoters commonly used to direct protein expression include the beta-lactamase (penicillinase), the lactose (lac), the tryptophan (Trp) (Goedel et al., Nucl. Acids Res. 8:4057, 1980), and the tac promoter systems, as well as the lambda-derived P.sub.L promoter and N-gene ribosome binding site (Simatake et al., Nature 292:128, 1981).

One particular bacterial expression system for P450 production is the *E. coli* pET expression system (Novagen). According to this expression system, DNA encoding a P450 is inserted into a pET vector in an orientation designed to allow expression. Since the P450 gene is under the control of the T7 regulatory signals, P450 expression is dependent on inducing the expression of T7 RNA polymerase in the host cell. This is typically achieved using host strains which express T7 RNA polymerase in response to IPTG induction. Once produced, recombinant P450 is then isolated according to standard methods known in the art, for example, those described herein.

Another bacterial expression system for P450 production is the pGEX expression system (Pharmacia). This system employs a GST gene fusion system that is designed for high-level expression of a gene or gene fragment as a fusion protein with rapid purification and recovery of the functional gene product. The P450 of interest is fused to the carboxyl terminus of the glutathione S-transferase protein from *Schistosoma japonicum* and is readily purified from bacterial lysates by affinity chromatography using Glutathione Sepharose 4B. Fusion proteins can be recovered under mild conditions by elution with glutathione. Cleavage of the glutathione S-transferase domain from the fusion protein is facilitated by the presence of recognition sites for site-specific proteases upstream of this domain. For example, proteins expressed in pGEX-2T plasmids may be cleaved with thrombin; those expressed in pGEX-3X may be cleaved with factor Xa.

Other prokaryotic systems useful for expressing eukaryotic P450s are described by Cooper (Mutat. Res. 454:45-52, 2000) and Dong et al. (Arch. Biochem. Biophys. 327:254-

259, 1996). In addition, strategies for enhancing the prokaryotic expression of a cytochrome P450 in combination with cytochrome reductase are described in Porter et al. (Drug. Metab. Rev. 31:159-174, 1999).

For eukaryotic expression, the method of transformation or transfection and the choice of vehicle for expression of the P450 will depend on the host system selected. Transformation and transfection methods of numerous organisms, for example, the baker's yeast *Saccharomyces cerevisiae*, are described, e.g., in Ausubel et al. (supra); Weissbach and Weissbach, Methods for Plant Molecular Biology, Academic Press, 1989; Gelvin et al., Plant Molecular Biology Manual, Kluwer Academic Publishers, 1990; Kindle, K., Proc. Natl. Acad. Sci. U.S.A. 87:1228 (1990); Potrykus, I., Annu. Rev. Plant Physiol. Plant Mol. Biology 42:205 (1991); and BioRad (Hercules, Calif.) Technical Bulletin #1687 (Biological Particle Delivery Systems). Expression vehicles may be chosen from those provided, e.g., in Cloning Vectors: A Laboratory Manual (P. H. Pouwels et al., 1985, Supp. 1987); Gasser and Fraley (supra); Clontech Molecular Biology Catalog (Catalog 1992/93 Tools for the Molecular Biologist, Palo Alto, Calif.); and the references cited above.

One preferred eukaryotic expression system is the mouse 3T3 fibroblast host cell transfected with a pMAMneo expression vector (Clontech). pMAMneo provides: an RSV-LTR enhancer linked to a dexamethasone-inducible MMTV-LTR promoter, an SV40 origin of replication which allows replication in mammalian systems, a selectable neomycin gene, and SV40 splicing and polyadenylation sites. DNA encoding a P450 is inserted into the pMAMneo vector in an orientation designed to allow expression. The recombinant P450 is then isolated as described below. Other preferable host cells which may be used in conjunction with the pMAMneo expression vehicle include COS cells and CHO cells (ATCC Accession Nos. CRL 1650 and CCL 61, respectively).

Alternatively, if desired, a P450 is produced by a stably-transfected mammalian cell line. A number of vectors suitable for stable transfection of mammalian cells are available to the public, e.g., see Pouwels et al. (supra); methods for constructing such cell lines are also publicly available, e.g., in Ausubel et al. (supra). In one example, cDNA encoding the P450 is cloned into an expression vector which includes the dihydrofolate reductase (DHFR) gene. Integration of the plasmid and, therefore, the P450-encoding gene into the host cell chromosome is selected for by inclusion of 0.01-300 μ M methotrexate in the cell culture medium (as described in Ausubel et al., supra). This dominant selection can be accomplished in most cell types. Recombinant protein expression can be increased by DHFR-mediated amplification of the transfected gene. Methods for selecting cell lines bearing gene amplifications are described in Ausubel et al. (supra); such methods generally involve extended culture in medium containing gradually increasing levels of methotrexate. DHFR-containing expression vectors commonly used for this purpose include pCVSEII-DHFR and pAdD26SV(A) (described in Ausubel et al., supra). Any of the host cells described above or, preferably, a DHFR-deficient CHO cell line (for example, CHO DHFR cells, ATCC Accession Number CRL 9096) are among the host cells preferred for DHFR selection of a stably-transfected cell line or DHFR-mediated gene amplification.

A cytochrome P450 may also be produced in insect cells, such cells include, without limitation, *Spodoptera frugiperda* Sf-21, or *Drosophila melanogaster* Schneider (SL-2) cells. For P450 production, insect cells are typically infected with a baculovirus, for example, *Autographa californica* Multiple Nuclear Polyhedrosis Virus (AcMNPV) containing an

expression cassette for such a protein, e.g., cytochrome P450, at a multiplicity of infection of 1 to 10. The infected cells are generally cultured in a standard insect cell culture medium for 24 to 48 hours prior to recovering the protein using standard molecular biology techniques. If desired, a P450 polypeptide may also be produced in insect cells directly transfected with a DNA construct containing an expression cassette encoding the P450.

Furthermore, any of the cytochrome P450s described herein may be produced in yeast, for example, *Pichia pastoris*. In order to produce the P450, yeast cells are transformed with an expression cassette containing, for example, a promoter such as the AOX1 or phosphoglycerate kinase gene promoter, the P450 gene to be expressed, and a terminator. Such an expression cassette may contain an origin of replication or it may be integrated into the yeast genomic DNA. The expression cassette is generally introduced by lithium acetate transformation or by the use of spheroplasts. In order to select for successfully transformed cells, the yeast are plated, for example, on minimal media which only allows yeast carrying the introduced expression cassette to grow.

In addition, expression of recombinant proteins in yeast using a *Hansenula polymorpha* expression system is described in U.S. Pat. Nos. 5,741,674 and 5,672,487.

A P450 may also be produced by a stably-transfected plant cell line or by a transgenic plant. Such genetically-engineered plants are useful for a variety of industrial and agricultural applications as discussed below. Importantly, this invention is applicable to gymnosperms and angiosperms, and will be readily applicable to any new or improved transformation or regeneration method.

A number of vectors suitable for stable transfection of plant cells or for the establishment of transgenic plants are available to the public; such vectors are described in Pouwels et al. (supra), Weissbach and Weissbach (supra), and Gelvin et al. (supra). Methods for constructing such cell lines are described in, e.g., Weissbach and Weissbach (supra), and Gelvin et al. (supra). Typically, plant expression vectors include (1) a cloned P450 gene under the transcriptional control of 5' and 3' regulatory sequences and (2) a dominant selectable marker. Such plant expression vectors may also contain, if desired, a promoter regulatory region (for example, one conferring inducible or constitutive expression, or environmentally- or developmentally-regulated, or pathogen- or wound-inducible, or cell- or tissue-specific expression), a transcription initiation start site, a ribosome binding site, an RNA processing signal, a transcription termination site, and/or a polyadenylation signal.

The P450 DNA sequence of the invention may, if desired, be combined with other DNA sequences in a variety of ways. The P450 DNA sequence of the invention may be employed with all or part of the gene sequences normally associated with a P450. In its component parts, a DNA sequence encoding a P450 is combined in a DNA construct having a transcription initiation control region capable of promoting transcription and translation in a host cell.

In general, the constructs will involve regulatory regions functional in plants which provide for production of a P450 as discussed herein. The open reading frame coding for the P450, or a functional fragment thereof, will be joined at its 5' end to a transcription initiation regulatory region such as the sequence naturally found in the 5' upstream region of a P450 structural gene, for example, a CYP71D20 (SEQ ID NO:2) or CYP71D21 (SEQ ID NO:4) gene. Numerous other transcription initiation regions are available which provide for constitutive or inducible regulation.

For applications when developmental, cell, tissue, hormonal, environmental, or pathogen-inducible expression are desired, appropriate 5' upstream non-coding regions are obtained from other genes; for example, from genes regulated during seed development, embryo development, leaf development, or in response to a pathogen.

Regulatory transcript termination regions may also be provided in DNA constructs of this invention as well. Transcript termination regions may be provided by the DNA sequence encoding a P450 or any convenient transcription termination region derived from a different gene source. The transcript termination region will contain preferably at least 1-3 kb of sequence 3' to the structural gene from which the termination region is derived.

An example of a useful plant promoter according to the invention is a caulimovirus promoter, such as, a cauliflower mosaic virus (CaMV) promoter. These promoters confer high levels of expression in most plant tissues, and the activity of these promoters is not dependent on virally encoded proteins. CaMV is a source for both the 35S and 19S promoters. In most tissues of transgenic plants, the CaMV 35S promoter is a strong promoter (see, e.g., Odell et al., *Nature* 313:810, 1985). The CaMV promoter is also highly active in monocots (see, e.g., Dekeyser et al., *Plant Cell* 2:591, 1990; Terada and Shimamoto, *Mol. Gen. Genet.* 220:389, 1990). Moreover, activity of this promoter can be further increased (i.e., between 2-10 fold) by duplication of the CaMV 35S promoter (see e.g., Kay et al., *Science* 236:1299, 1987; Ow et al., *Proc. Natl. Acad. Sci. U.S.A.* 84:4870, 1987; and Fang et al., *Plant Cell* 1:141, 1989). Other useful plant promoters include, without limitation, the nopaline synthase promoter (An et al., *Plant Physiol.* 88:547, 1988) and the octopine synthase promoter (Fromm et al., *Plant Cell* 1:977, 1989).

For certain applications, it may be desirable to produce the P450 gene product in an appropriate tissue, at an appropriate level, or at an appropriate developmental time. For this purpose, there is an assortment of gene promoters, each with its own distinct characteristics embodied in its regulatory sequences, which have been shown to be regulated in response to the environment, hormones, and/or developmental cues. These include gene promoters that are responsible for heat-regulated gene expression (see, e.g., Callis et al., *Plant Physiol.* 88:965, 1988; Takahashi and Komeda, *Mol. Gen. Genet.* 219:365, 1989; and Takahashi et al., *Plant J.* 2:751, 1992); light-regulated gene expression (e.g., the pea *rbcs-3A* described by Kuhlemeier et al. (*Plant Cell* 1:471, 1989); the maize *rbcs* promoter described by Schaffner and Sheen (*Plant Cell* 3:997, 1991); or the chlorophyll *a/b*-binding protein gene found in pea described by Simpson et al. (*EMBO J.* 4:2723, 1985)); hormone-regulated gene expression (for example, the abscisic acid (ABA) responsive sequences from the *Em* gene of wheat described by Marcotte et al. (*Plant Cell* 1:969, 1989); the ABA-inducible *HVA1* and *HVA22*, and the *rd29A* promoters described for barley and *Arabidopsis* by Straub et al. (*Plant Cell* 6:617, 1994), Shen et al. (*Plant Cell* 7:295, 1994)); and wound-induced gene expression (for example, of wu1 described by Siebertz et al. (*Plant Cell* 1:961, 1989); or organ-specific gene expression (for example, of the tuber-specific storage protein gene described by Roshal et al. (*EMBO J.* 6: 1155, 1987); the 23-kDa zein gene from maize described by Schernthaner et al. (*EMBO J.* 7:1249, 1988); or the French bean beta-phaseolin gene described by Bustos et al. (*Plant Cell* 1:839, 1989); and pathogen-inducible gene expression described by Chappell et al. in U.S. Ser. Nos. 08/471,983; 08/443,639; and 08/577,483; hereby incorporated by reference.

Plant expression vectors may also optionally include RNA processing signals, for example, introns, which have been shown to be important for efficient RNA synthesis and accumulation (Callis et al., *Genes and Dev.* 1:1183, 1987). The location of the RNA splice sequences can dramatically influence the level of transgene expression in plants. In view of this fact, an intron may be positioned upstream or downstream of a P450-encoding sequence in the transgene to modulate levels of gene expression.

In addition to the aforementioned 5' regulatory control sequences, the expression vectors may also include regulatory control regions which are generally present in the 3' regions of plant genes (Thornburg et al., *Proc. Natl. Acad. Sci. U.S.A.* 84:744, 1987; An et al., *Plant Cell* 1:115, 1989). For example, the 3' terminator region may be included in the expression vector to increase stability of the mRNA. One such terminator region may be derived from the PI-II terminator region of potato. In addition, other commonly used terminators are derived from the octopine or nopaline synthase signals.

The plant expression vector also typically contains a dominant selectable marker gene used to identify those cells that have become transformed. Useful selectable genes for plant systems include genes encoding antibiotic resistance genes, for example, those encoding resistance to hygromycin, kanamycin, bleomycin, G418, streptomycin, or spectinomycin. Genes required for photosynthesis may also be used as selectable markers in photosynthetic-deficient strains. Alternatively, the green-fluorescent protein from the jellyfish *Aequorea victoria* may be used as a selectable marker (Sheen et al., *Plant J.* 8:777, 1995; Chiu et al., *Current Biology* 6:325, 1996). Finally, genes encoding herbicide resistance may be used as selectable markers; useful herbicide resistance genes include the bar gene encoding the enzyme phosphinothricin acetyltransferase and conferring resistance to the broad-spectrum herbicide BASTA (Hoechst AG, Frankfurt, Germany).

Efficient use of selectable markers is facilitated by a determination of the susceptibility of a plant cell to a particular selectable agent and a determination of the concentration of this agent which effectively kills most, if not all, of the transformed cells. Some useful concentrations of antibiotics for tobacco transformation include, e.g., 75-100 µg/mL (kanamycin), 20-50 µg/mL (hygromycin), or 5-10 µg/mL (bleomycin). A useful strategy for selection of transformants for herbicide resistance is described, e.g., by Vasil et al., *supra*.

It should be readily apparent to one skilled in the art of molecular biology, especially in the field of plant molecular biology, that the level of gene expression is dependent, not only on the combination of promoters, RNA processing signals, and terminator elements, but also on how these elements are used to increase the levels of selectable marker gene expression.

Plant Transformation

Upon construction of the plant expression vector, several standard methods are available for introduction of the vector into a plant host, thereby generating a transgenic plant. These methods include (1) *Agrobacterium*-mediated transformation (*A. tumefaciens* or *A. rhizogenes*) (see, e.g., Lichtenstein and Fuller, In: *Genetic Engineering*, vol. 6, PWJ Rigby, ed, London, Academic Press, 1987; and Lichtenstein, C. P., and Draper, J., In: *DNA Cloning*, Vol II, D. M. Glover, ed, Oxford, IRI Press, 1985); (2) the particle delivery system (see, e.g., Gordon-Kamm et al., *Plant Cell* 2:603, 1990; or BioRad Technical Bulletin 1687, *supra*); (3) microinjection protocols (see, e.g., Green et al., *supra*); (4) polyethylene glycol (PEG) procedures (see, e.g., Draper et al., *Plant Cell Physiol.* 23:451, 1982; or e.g., Zhang and Wu, *Theor. Appl. Genet.*

76:835, 1988); (5) liposome-mediated DNA uptake (see, e.g., Freeman et al., Plant Cell Physiol. 25:1353, 1984); (6) electroporation protocols (see, e.g., Gelvin et al., supra; Dekeyser et al., supra; Fromm et al., Nature 319:791, 1986; Sheen, Plant Cell 2:1027, 1990; or Jang and Sheen, Plant Cell 6:1665, 1994); and (7) the vortexing method (see, e.g., Kindle, supra). The method of transformation is not critical to the present invention. Any method which provides for efficient transformation may be employed. As newer methods are available to transform crops or other host cells, they may be directly applied.

The following is an example outlining one particular technique, an *Agrobacterium*-mediated plant transformation. By this technique, the general process for manipulating genes to be transferred into the genome of plant cells is carried out in two phases. First, cloning and DNA modification steps are carried out in *E. coli*, and the plasmid containing the gene construct of interest is transferred by conjugation or electroporation into *Agrobacterium*. Second, the resulting *Agrobacterium* strain is used to transform plant cells. Thus, for the generalized plant expression vector, the plasmid contains an origin of replication that allows it to replicate in *Agrobacterium* and a high copy number origin of replication functional in *E. coli*. This permits facile production and testing of transgenes in *E. coli* prior to transfer to *Agrobacterium* for subsequent introduction into plants. Resistance genes can be carried on the vector, one for selection in bacteria, for example, streptomycin, and another that will function in plants, for example, a gene encoding kanamycin resistance or herbicide resistance. Also present on the vector are restriction endonuclease sites for the addition of one or more transgenes and directional T-DNA border sequences which, when recognized by the transfer functions of *Agrobacterium*, delimit the DNA region that will be transferred to the plant.

In another example, plant cells may be transformed by shooting into the cell tungsten microprojectiles on which cloned DNA is precipitated. In the Biolistic Apparatus (Bio-Rad) used for the shooting, a gunpowder charge (22 caliber Power Piston Tool Charge) or an air-driven blast drives a plastic macroprojectile through a gun barrel. An aliquot of a suspension of tungsten particles on which DNA has been precipitated is placed on the front of the plastic macroprojectile. The latter is fired at an acrylic stopping plate that has a hole through it that is too small for the macroprojectile to pass through. As a result, the plastic macroprojectile smashes against the stopping plate, and the tungsten microprojectiles continue toward their target through the hole in the plate. For the present invention, the target can be any plant cell, tissue, seed, or embryo. The DNA introduced into the cell on the microprojectiles becomes integrated into either the nucleus or the chloroplast.

In general, transfer and expression of transgenes in plant cells are now routine practices to those skilled in the art, and have become major tools to carry out gene expression studies in plants and to produce improved plant varieties of agricultural or commercial interest.

Transgenic Plant Regeneration

Plants cells transformed with plant expression vectors can be regenerated, for example, from single cells, callus tissue, or leaf discs according to standard plant tissue culture techniques. It is well known in the art that various cells, tissues, and organs from almost any plant can be successfully cultured to regenerate an entire plant; such techniques are described, e.g., in Vasil (supra), Green et al. (supra), Weissbach and Weissbach (supra) and Gelvin et al. (supra).

In one particular example, a cloned P450, under the control of the EAS4 promoter and the nopaline synthase terminator

and carrying a selectable marker (for example, kanamycin resistance), is transformed into *Agrobacterium*. Transformation of leaf discs (for example, of tobacco leaf discs), with vector-containing *Agrobacterium* is carried out as described by Horsch et al. (Science 227:1229, 1985). Putative transformants are selected after a few weeks (for example, 3 to 5 weeks) on plant tissue culture media containing kanamycin (e.g., 100 µg/mL). Kanamycin-resistant shoots are then placed on plant tissue culture media without hormones for root initiation. Kanamycin-resistant plants are then selected for greenhouse growth. If desired, seeds from self-fertilized transgenic plants can then be sowed in soil-less medium and grown in a greenhouse. Kanamycin-resistant progeny are selected by sowing surface sterilized seeds on hormone-free kanamycin-containing media. Analysis for the integration of the transgene is accomplished by standard techniques (see, for example, Ausubel et al. (supra); Gelvin et al. (supra)).

Transgenic plants expressing the selectable marker are then screened for transmission of the transgene DNA by standard immunoblot and DNA detection techniques. Each positive transgenic plant and its transgenic progeny is unique in comparison to other transgenic plants established with the same transgene. Integration of the transgene DNA into the plant genomic DNA is in most cases random, and the site of integration can profoundly affect the levels and the tissue and developmental patterns of transgene expression. Consequently, a number of transgenic lines are usually screened for each transgene to identify and select plants with the most appropriate expression profiles.

Transgenic lines are generally evaluated for levels of transgene expression. Expression at the RNA level is determined initially to identify and quantitate expression-positive plants. Standard techniques for RNA analysis are employed and include PCR amplification assays using oligonucleotide primers designed to amplify only transgene RNA templates and solution hybridization assays using transgene-specific probes (see, e.g., Ausubel et al. (supra)). The RNA-positive plants are then analyzed for protein expression by Western immunoblot analysis using specific antibodies to the P450 (see, e.g., Ausubel et al., supra). In addition, in situ hybridization and immunocytochemistry according to standard protocols can be done using transgene-specific nucleotide probes and antibodies, respectively, to localize sites of expression within transgenic tissue.

Once the recombinant P450 is expressed in any cell or in a transgenic plant (for example, as described above), it may be isolated, e.g., using affinity chromatography. In one example, an anti-P450 antibody (e.g., produced as described in Ausubel et al., supra, or by any standard technique) may be attached to a column and used to isolate the polypeptide. Lysis and fractionation of P450-producing cells prior to affinity chromatography may be performed by standard methods (see, e.g., Ausubel et al., supra). Once isolated, the recombinant protein can, if desired, be further purified, for example, by high performance liquid chromatography (see, e.g.,

Fisher, Laboratory Techniques in Biochemistry and Molecular Biology, eds., Work and Burdon, Elsevier, 1980).

These general techniques of polypeptide expression and purification can also be used to produce and isolate useful P450 fragments or analogs.

Use

The aforementioned cytochrome P450 polypeptides of the invention are useful in the biosynthesis of hormones, lipids, and secondary metabolites, and may also help plants tolerate potentially harmful exogenous chemicals such as herbicides, pesticides, and pollutants. In addition, such cytochrome P450

polypeptides are useful in the chemical defense of plants against insects, as well as against bacterial, viral, and fungal infection.

Engineering Plant Disease Resistance

Plasmid constructs designed for the expression of a P450 gene product are useful, for example, for activating plant defense pathways that confer anti-pathogenic properties to a transgenic plant, for example, the production of phytoalexins. P450 genes that are isolated from a host plant (e.g., *Nicotiana*) may be engineered for expression in the same plant, a closely related species, or a distantly related plant species. For example, a P450 gene may be engineered for constitutive low-level expression and then transformed into a *Nicotiana* host plant. Alternatively, the P450 gene may be engineered for expression in other solanaceous plants, including, but not limited to, potato and tomato. To achieve pathogen resistance, it is important to express a P450 protein at an effective level. Evaluation of the level of pathogen protection conferred to a plant by ectopic expression of the P450 gene is determined according to conventional methods and assays.

Industrial Applications

The invention also includes engineering host cells to include novel isoprenoid metabolic pathways useful in the production of new isoprenoid compounds. By introducing genes encoding an isoprenoid synthase (as disclosed in U.S. Pat. No. 5,824,774 and WO 00/17327) and a cytochrome P450, an acetyltransferase, a methyl transferase, a fatty acyltransferase, or a combination thereof, various isoprenoid reaction products may be modified, controlled, or manipulated, resulting in enhancement of production of numerous isoprenoid reaction products, for example, the production of novel monoterpenes, diterpenes, and sesquiterpenes. Such compounds are useful as phytoalexins, insecticides, perfumes, and pharmaceuticals such as anti-bacterial and fungal agents.

In one working example, an isoprenoid synthase or a chimeric isoprenoid synthase (as disclosed in U.S. Pat. No. 5,824,774 and WO 00/17327) and a P450 gene are introduced

into yeast, for example, using any of the procedures described herein. If desired, such cells may also express, either independently or in combination, an acetyltransferase (see, for example, Walker et al., Proc. Natl. Acad. Sci. U.S.A. 18:583-587, 2000), a methylase transferase gene (see, for example, Diener et al., Plant Cell 12:853-870, 2000), or a fatty acyltransferase gene, as well as a cytochrome reductase. Cells are then cultured under standard conditions and the production of isoprenoid compounds is assayed according to methods known in the art. Isoprenoid compounds are further purified according to methods well known in the art. Cells expressing novel isoprenoid compounds are taken as useful in the invention.

Such methods provide a unique approach for producing novel isoprenoid starting materials and end products. Either prokaryotic or eukaryotic cells transformed with any of the aforementioned enzymes (or combinations thereof) may be used. Moreover, isoprenoid compounds may be produced in any number of ways known in the art including an in vitro combination of purified enzymes with an appropriate substrate or direct fermentation using a host cell which expresses any combination of the aforementioned enzymes and the appropriate substrates sufficient to drive production of isoprenoid compounds.

The invention is also useful for the production of insect attractants and deterrents, which may either deter insect pests or attract insect predators. In addition, the invention is also useful for generating novel flavorings and perfumes.

Other Embodiments

From the foregoing description, one skilled in the art can easily ascertain the essential characteristics of this invention, and can make various changes and modifications of the invention to adapt it to various usages and conditions. Thus, other embodiments are also within the claims.

All publications and patents mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication or patent was specifically and individually indicated to be incorporated by reference.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 43

<210> SEQ ID NO 1

<211> LENGTH: 473

<212> TYPE: PRT

<213> ORGANISM: *Nicotiana tabacum*

<400> SEQUENCE: 1

```
Met Gln Phe Phe Ser Leu Val Ser Ile Phe Leu Phe Leu Ala Phe Leu
1           5           10           15

Phe Leu Leu Arg Lys Trp Lys Asn Ser Asn Ser Gln Ser Lys Lys Leu
20           25           30

Pro Pro Gly Pro Trp Lys Ile Pro Ile Leu Gly Ser Met Leu His Met
35           40           45

Ile Gly Gly Glu Pro His His Val Leu Arg Asp Leu Ala Lys Lys Tyr
50           55           60

Gly Pro Leu Met His Leu Gln Leu Gly Glu Ile Ser Ala Val Val Val
65           70           75           80

Thr Ser Arg Asp Met Ala Lys Glu Val Leu Lys Thr His Asp Val Val
85           90           95

Phe Ala Ser Arg Pro Lys Ile Val Ala Met Asp Ile Ile Cys Tyr Asn
100          105          110
```

-continued

Gln Ser Asp Ile Ala Phe Ser Pro Tyr Gly Asp His Trp Arg Gln Met
 115 120 125
 Arg Lys Ile Cys Val Met Glu Leu Leu Asn Ala Lys Asn Val Arg Ser
 130 135 140
 Phe Ser Ser Ile Arg Arg Asp Glu Val Val Arg Leu Ile Asp Ser Ile
 145 150 155 160
 Arg Ser Asp Ser Ser Ser Gly Glu Leu Val Asn Phe Thr Gln Arg Ile
 165 170 175
 Ile Trp Phe Ala Ser Ser Met Thr Cys Arg Ser Ala Phe Gly Gln Val
 180 185 190
 Leu Lys Gly Gln Asp Ile Phe Ala Lys Lys Ile Arg Glu Val Ile Gly
 195 200 205
 Leu Ala Glu Gly Phe Asp Val Val Asp Ile Phe Pro Thr Tyr Lys Phe
 210 215 220
 Leu His Val Leu Ser Gly Met Lys Arg Lys Leu Leu Asn Ala His Leu
 225 230 235 240
 Lys Val Asp Ala Ile Val Glu Asp Val Ile Asn Glu His Lys Lys Asn
 245 250 255
 Leu Ala Ala Gly Lys Ser Asn Gly Ala Leu Glu Asp Met Phe Ala Ala
 260 265 270
 Gly Thr Glu Thr Ser Ser Thr Thr Thr Val Trp Ala Met Ala Glu Met
 275 280 285
 Met Lys Asn Pro Ser Val Phe Thr Lys Ala Gln Ala Glu Val Arg Glu
 290 295 300
 Ala Phe Arg Asp Lys Val Ser Phe Asp Glu Asn Asp Val Glu Glu Leu
 305 310 315 320
 Lys Tyr Leu Lys Leu Val Ile Lys Glu Thr Leu Arg Leu His Pro Pro
 325 330 335
 Ser Pro Leu Leu Val Pro Arg Glu Cys Arg Glu Asp Thr Asp Ile Asn
 340 345 350
 Gly Tyr Thr Ile Pro Ala Lys Thr Lys Val Met Val Asn Val Trp Ala
 355 360 365
 Leu Gly Arg Asp Pro Lys Tyr Trp Asp Asp Ala Glu Ser Phe Lys Pro
 370 375 380
 Glu Arg Phe Glu Gln Cys Ser Val Asp Phe Phe Gly Asn Asn Phe Glu
 385 390 395 400
 Phe Leu Pro Phe Gly Gly Gly Arg Arg Ile Cys Pro Gly Met Ser Phe
 405 410 415
 Gly Leu Ala Asn Leu Tyr Leu Pro Leu Ala Gln Leu Leu Tyr His Phe
 420 425 430
 Asp Trp Lys Leu Pro Thr Gly Ile Met Pro Arg Asp Leu Asp Leu Thr
 435 440 445
 Glu Leu Ser Gly Ile Thr Ile Ala Arg Lys Gly Asp Leu Tyr Leu Asn
 450 455 460
 Ala Thr Pro Tyr Gln Pro Ser Arg Glu
 465 470

<210> SEQ ID NO 2

<211> LENGTH: 1660

<212> TYPE: DNA

<213> ORGANISM: Nicotiana tabacum

<400> SEQUENCE: 2

ggatggtcta ataatactcc atttatctcc gaaaatgcaa ttcttcagct tggtttccat

60

-continued

tttctcttcc	ctagctttcc	tatttttggt	gaggaaatgg	agaactcca	atagccaaag	120
caaaaaattg	ccaccaggtc	catggaaaat	accaatacta	ggaagtatgc	ttcatatgat	180
tggtggagaa	cgcaccatg	tccttagaga	tttagccaaa	aatatggac	cacttatgca	240
ccttcagtta	ggtgaaattt	ctgcagttgt	ggttacttct	agggacatgg	caaaagaagt	300
gctaaaaact	catgacgtcg	tttttgcatc	taggectaaa	attgtagcca	tggacattat	360
ctgttataac	cagtcgcaca	ttgcctttag	cccttatggc	gaccactgga	gacaaatgcg	420
taaaatttgt	gtcatggaac	ttotcaatgc	aaagaatggt	cggttcttca	gctccatcag	480
acgtgatgaa	gtcgttcgtc	tcattgactc	tatccggtea	gattcttctt	caggtgagct	540
agttaatttt	acgcagagga	tcatttggtt	tgcaagctcc	atgacgtgta	gatcagcatt	600
tgggcaagta	ctcaaggggc	aagacatatt	tgccaaaaag	atcagagaag	taataggatt	660
agcagaaggc	tttgatgtgg	tagacatctt	ccctacatac	aagtttcttc	atgttctcag	720
tgggatgaag	cgtaaacttt	tgaatgccca	ccttaaggta	gacgccattg	ttgaggatgt	780
catcaacgag	cacaagaaaa	atcttgacgc	tggcaaaaag	aatggcgcat	tagaggacat	840
gtttgctgcc	ggaacagaaa	cttcatcaac	aacaactgta	tgggctatgg	ctgaaatgat	900
gaagaatcca	agtgtattca	ccaaagctca	agcagaagtg	cgagaagcct	ttagggacaa	960
agtatctttt	gatgaaaatg	atgtggagga	gctgaaatac	ttaaagttag	tcattaaaga	1020
aactttgaga	cttcatccac	cgtctccact	tttggcccc	agagaatgca	gggaagatac	1080
ggatataaac	ggctacacta	ttcctgcaaa	gaccaaagtt	atggttaatg	ttgggcatt	1140
gggaagagat	ccaaaatatt	gggatgacgc	ggaaagcttt	aagccagaga	gatttgagca	1200
atgctctgtg	gatttttttg	gtaataattt	tgagtttctt	ccctttggcg	gtggacggag	1260
aatttgctct	ggaatgtcat	ttggttttag	taatctttac	ttgccattgg	ctcaattact	1320
ctatcacttt	gactggaaac	tcccaaccgg	aatcatgcc	agagacttag	acttgaccga	1380
attatcggga	ataactattg	ctagaaaggg	tgacctttac	ttaaatgcc	ctccttatca	1440
accttctcga	gagtaattca	atattggcat	aaacatttta	aatttccttc	atcaacctca	1500
atattgtaca	ataatcatc	ttctgggtgt	ataggcttta	tcgatttcca	atacatgtat	1560
tctttattaa	aaaatgtatc	acattccatg	tagaaggagg	acgcaccaat	taattgtgcc	1620
atgattttag	ggtaacttgt	tccatcttaa	aaaaaaaaa			1660

<210> SEQ ID NO 3

<211> LENGTH: 474

<212> TYPE: PRT

<213> ORGANISM: Nicotiana tabacum

<400> SEQUENCE: 3

Met	Glu	Phe	Phe	Ser	Leu	Val	Ser	Ile	Phe	Leu	Phe	Leu	Ser	Phe	Leu
1				5					10					15	
Phe	Leu	Leu	Arg	Lys	Cys	Lys	Asn	Ser	Asn	Ser	Gln	Thr	Lys	Gln	Leu
			20				25						30		
Pro	Pro	Gly	Pro	Trp	Lys	Ile	Pro	Ile	Leu	Gly	Ser	Met	Leu	His	Met
		35				40					45				
Leu	Gly	Gly	Glu	Pro	His	His	Ile	Leu	Arg	Asp	Leu	Ala	Lys	Lys	Tyr
	50				55					60					
Gly	Pro	Ile	Met	His	Leu	Gln	Phe	Gly	Glu	Ile	Ser	Ala	Val	Val	Val
65				70				75					80		
Thr	Ser	Arg	Glu	Met	Ala	Lys	Glu	Val	Leu	Lys	Thr	His	Asp	Val	Val
			85					90					95		

-continued

Phe	Ala	Ser	Arg	Pro	Lys	Ile	Val	Ala	Met	Asp	Ile	Ile	Cys	Tyr	Asn	
			100					105					110			
Gln	Ser	Asp	Ile	Ala	Phe	Ser	Pro	Tyr	Gly	Asp	His	Trp	Arg	Gln	Met	
		115					120					125				
Arg	Lys	Ile	Cys	Val	Met	Glu	Leu	Leu	Asn	Ala	Lys	Asn	Val	Arg	Ser	
	130					135					140					
Phe	Ser	Ser	Ile	Arg	Arg	Asp	Glu	Val	Val	Arg	Leu	Ile	Asp	Ser	Ile	
145					150					155					160	
Arg	Ser	Asp	Ser	Ser	Ser	Gly	Glu	Leu	Val	Asn	Phe	Thr	Gln	Arg	Ile	
			165						170					175		
Ile	Trp	Phe	Ala	Ser	Ser	Met	Thr	Cys	Arg	Ser	Ala	Phe	Gly	Gln	Val	
		180						185					190			
Leu	Lys	Gly	Gln	Asp	Val	Phe	Ala	Lys	Lys	Ile	Arg	Glu	Val	Ile	Gly	
	195						200					205				
Leu	Ala	Glu	Gly	Phe	Asp	Val	Ala	Asp	Ile	Phe	Pro	Ser	Tyr	Lys	Phe	
	210					215					220					
Leu	His	Val	Leu	Ser	Gly	Met	Lys	Arg	Lys	Leu	Leu	Asn	Ala	His	Leu	
225					230					235					240	
Lys	Val	Asp	Ala	Ile	Val	Glu	Asp	Val	Ile	Asn	Glu	His	Lys	Lys	Asn	
			245						250					255		
Leu	Ala	Thr	Gly	Lys	Thr	Asn	Gly	Ala	Leu	Gly	Glu	Asp	Met	Phe	Ala	
		260					265						270			
Ala	Gly	Thr	Glu	Thr	Ser	Ser	Thr	Thr	Thr	Val	Trp	Ala	Met	Ala	Glu	
	275						280					285				
Met	Met	Lys	Asn	Pro	Asn	Val	Phe	Asn	Lys	Ala	Gln	Ala	Glu	Val	Arg	
	290					295					300					
Glu	Thr	Phe	Lys	Asp	Lys	Val	Thr	Phe	Asp	Glu	Ile	Asp	Ala	Glu	Glu	
305					310					315					320	
Leu	Glu	Tyr	Leu	Lys	Leu	Val	Ile	Lys	Glu	Thr	Leu	Arg	Leu	His	Pro	
			325						330					335		
Pro	Ser	Pro	Leu	Leu	Val	Pro	Arg	Glu	Cys	Arg	Glu	Asp	Thr	Asp	Ile	
		340						345					350			
Asn	Gly	Tyr	Thr	Ile	Pro	Ala	Lys	Thr	Lys	Val	Met	Val	Asn	Val	Trp	
	355						360					365				
Ala	Leu	Gly	Arg	Asp	Pro	Lys	Tyr	Trp	Asp	Asp	Ala	Glu	Ser	Phe	Lys	
	370					375					380					
Pro	Glu	Arg	Phe	Glu	Gln	Cys	Ser	Val	Asp	Phe	Phe	Gly	Asn	Asn	Phe	
385					390					395					400	
Glu	Phe	Leu	Pro	Phe	Gly	Gly	Gly	Arg	Arg	Ile	Cys	Pro	Gly	Met	Ser	
			405						410					415		
Phe	Gly	Leu	Ala	Asn	Leu	Tyr	Leu	Pro	Leu	Ala	Gln	Leu	Leu	Tyr	His	
		420					425						430			
Phe	Asp	Trp	Lys	Leu	Pro	Ser	Gly	Met	Met	Pro	Gly	Asp	Leu	Asp	Leu	
	435						440					445				
Thr	Glu	Leu	Ala	Gly	Ile	Thr	Ile	Ala	Arg	Lys	Gly	Asp	Leu	Tyr	Leu	
	450					455					460					
Met	Ala	Thr	Pro	Tyr	Gln	Pro	Ser	Arg	Glu							
465					470											

<210> SEQ ID NO 4
 <211> LENGTH: 1614
 <212> TYPE: DNA
 <213> ORGANISM: *Nicotiana tabacum*
 <400> SEQUENCE: 4

-continued

```

ggatggtcta ataatectcc atttatctcc caaaatggaa ttcttcagct tggtttccat    60
attcctattc ctatctttcc tctttttggt aaggaaatgt aagaactcca atagccaaac    120
caaaccaattg cctccaggtc catggaaaat accaatacta ggaagtatgc ttcatatgct    180
tgggtggagaa ccacaccata tccttaggga tttagccaaa aaatatggac caattatgca    240
ccttcagttt ggtgaaattt ctgcagttgt ggttacttct agggagatgg caaaagaagt    300
gctaaaaact catgacgtag tttttgcatc taggcctaaa attgtggcca tggacattat    360
ctgttataac cagtctgata tcgcctttag cccttatggc gatcactgga gacaaatgcg    420
taaaatttgt gtcatggaac ttcttaatgc aaagaatggt cggtctttca gctcgatcag    480
acgtgatgaa gtcgttcgtc tcattgactc tattegatca gattcttctt ctggtgagct    540
agttaatttt acgcaaagga tcatttggtt cgcgagctcc atgacgtgta gatcagcatt    600
tgggcaagta cttaaagggc aagacgtatt tgccaaaaag attagagaag taataggggt    660
agcagaaggc tttgatgtgg ccgatatctt cccttcatac aagtttcttc atgttctcag    720
tggaatgaag cgtaaacctc tgaatgccca ccttaaggta gatgccattg ttgaggatgt    780
catcaacgag cacaagaaaa atcttgcaac tgggaaaact aatggagcat taggagacat    840
gtttgctgcc ggaacagaaa cttcatcaac aacaactgta tgggctatgg ctgaaatgat    900
gaagaatcca aatgtattca acaaagctca ggcagaagtg agagaaacct ttaaagacaa    960
agtaacattt gatgaaattg atgcagagga gctggaatac ttaaagttag ttattaaaga   1020
aactttgaga cttcatccac cgtctccact tttggtccca agagaatgta ggaagatac   1080
agatattaac ggctacacta ttcctgcgaa gaccaaagtt atggttaatg tttgggcatt   1140
gggaagagat ccaaaatatt gggatgacgc agaaagcttt aagccagaga gatttgagca   1200
atgctctgtg gatttttttg gtaataattt tgagtttctt ccctttggcg gtggacggag   1260
aatatgtcct ggtagtcat ttggttttag taatctttac ttgccattgg ctcaattgct   1320
atatcacttt gattggaaaac tcccagacgg aatgatgccc ggagacttgg acttgactga   1380
attagctgga ataacaattg ctagaaaggg tgacctttac ttaatggcta ctccttatca   1440
accttctcgc gaataattta atggcatcag gttttttaat tccattgtca acctcactat   1500
tgtacaagct ttctgatggt tcaggttttg ccgatttgta ataaatgtag tttttataat   1560
atgtatcata cccatgtaga agagggacga ttaattagtt gtaaaaaaaaa aaaa       1614

```

<210> SEQ ID NO 5

<211> LENGTH: 537

<212> TYPE: PRT

<213> ORGANISM: Nicotiana tabacum

<400> SEQUENCE: 5

```

Met Lys Asn Met Ala Lys Leu Leu Asn Lys Thr Ile Phe Cys Ile Leu
 1             5             10            15
Phe Thr Ile Ala Phe Leu Ser Phe Ala Lys Leu Leu Ser Ser Tyr Leu
      20            25            30
Ser Met Pro Phe Pro Leu Lys Tyr Met Ser Leu Ile Val Pro Leu Leu
      35            40            45
Pro Leu Ile Ile Asn Phe Leu Tyr Val Lys Pro Gln Asn Asn Leu Pro
      50            55            60
Pro Gly Pro Thr Ala Val Pro Ile Phe Gly Asn Trp Leu Gln Val Gly
      65            70            75            80
Asn Asp Leu Asn His Gln Leu Leu Ala Thr Met Ser Gln Thr Tyr Gly
      85            90            95

```

Pro 100	Ile	Phe	Leu	Leu	Lys	Leu	Gly	Ser	Lys	Asn	Leu	Ala	Val	Val	Ser
Asn 115	Pro	Glu	Leu	Ala	Asp	Gln	Val	Leu	His	Thr	Gln	Gly	Val	Glu	Phe
Gly 130	Ser	Arg	Pro	Arg	Asn	Val	Val	Phe	Asp	Ile	Phe	Thr	Gly	Asn	Gly
Gln 145	Asp	Met	Val	Phe	Thr	Ile	Tyr	Gly	Asp	His	Trp	Arg	Lys	Met	Arg
Arg	Ile	Met	Thr	Leu	Pro	Phe	Phe	Thr	Asn	Lys	Val	Val	His	Gln	Tyr
Ser	Asp	Met	Trp	Glu	Asn	Glu	Met	Asp	Leu	Val	Val	Asn	Asp	Leu	Lys
Lys	Asn	Glu	Lys	Val	Lys	Tyr	Glu	Gly	Ile	Val	Ile	Arg	Lys	Arg	Leu
Gln 210	Leu	Met	Leu	Tyr	Asn	Ile	Met	Tyr	Arg	Met	Met	Phe	Asp	Ala	Lys
Phe 225	Glu	Ser	Gln	Asn	Asp	Pro	Leu	Phe	Ile	Glu	Ala	Thr	Lys	Phe	Asn
Ser	Glu	Arg	Ser	Arg	Leu	Ala	Gln	Ser	Phe	Asp	Tyr	Asn	Tyr	Gly	Asp
Phe	Ile	Pro	Leu	Leu	Arg	Pro	Phe	Leu	Arg	Gly	Tyr	Leu	Asn	Lys	Cys
Lys	Asp	Leu	Gln	Thr	Arg	Arg	Leu	Ala	Phe	Phe	Asn	Asn	Tyr	Phe	Val
Glu 290	Lys	Arg	Arg	Lys	Ile	Met	Asp	Glu	Asn	Gly	Glu	Lys	His	Lys	Ile
Ser 305	Cys	Ala	Ile	Asp	His	Ile	Ile	Asp	Ala	Glu	Met	Lys	Gly	Glu	Ile
Asn	Glu	Gln	Asn	Val	Leu	Tyr	Ile	Val	Glu	Asn	Ile	Asn	Val	Ala	Ala
Ile	Glu	Thr	Thr	Leu	Trp	Ser	Met	Glu	Trp	Ala	Ile	Ala	Glu	Leu	Val
Asn 355	His	Pro	Ile	Val	Gln	Gln	Lys	Ile	Arg	Asp	Glu	Ile	Ser	Thr	Val
Leu 370	Lys	Gly	Arg	Ser	Val	Thr	Glu	Ser	Asn	Leu	His	Glu	Leu	Pro	Tyr
Leu 385	Leu	Ala	Thr	Val	Asn	Glu	Thr	Leu	Arg	Leu	His	Thr	Pro	Ile	Pro
Leu	Leu	Val	Pro	His	Met	Asn	Leu	Glu	Glu	Ala	Lys	Leu	Gly	Gly	Tyr
Thr	Ile	Pro	Lys	Glu	Thr	Lys	Val	Val	Val	Asn	Ala	Trp	Trp	Leu	Ala
Asn	Asn	Pro	Ala	Trp	Trp	Lys	Asn	Pro	Asn	Glu	Phe	Arg	Pro	Glu	Arg
Phe 450	Leu	Glu	Glu	Asp	Ser	Ser	Thr	Glu	Ala	Ala	Val	Ala	Gly	Gly	Lys
Val 465	Asp	Phe	Arg	Tyr	Leu	Pro	Phe	Gly	Met	Gly	Arg	Arg	Ser	Cys	Pro
Gly	Ile	Ile	Leu	Ala	Leu	Pro	Ile	Leu	Gly	Leu	Val	Ile	Ala	Lys	Leu
Val	Ser	Asn	Phe	Glu	Met	Gln	Gly	Pro	Gly	Val	Glu	Lys	Val	Asp	
Thr	Ser	Glu	Arg	Gly	Gly	Gln	Phe	Ser	Leu	His	Ile	Ala	Lys	His	Ser

-continued

Thr Val Val Phe Lys Pro Ile Ala Ala
530 535

<210> SEQ ID NO 6
<211> LENGTH: 1745
<212> TYPE: DNA
<213> ORGANISM: *Nicotiana tabacum*

<400> SEQUENCE: 6

```
cctctagcta atgaaaaaca tggccaaact tctcaacaag accatctttt gcattctctt    60
tacaattgca tttctttcat ttgccaaagt actgtcctcc tacctatcta tgcctttccc    120
tcttaagtac atgtcactta ttgtcccttt acttcccctt ataatcaact tcctctatgt    180
taagccccaa aacaacctcc cacctgggtcc aacagcagtc ccaatatattg gtaattggct    240
tcaagttggc aatgacttga accatcaact ccttgccacc atgtcacaaa cctacgggtcc    300
tatattttta ctcaaaacttg gttcaaaaaa cctagctgtg gtatcaaac cagagctagc    360
tgaccaagtt ctacacacac aaggggtcga gtttgggtcc cgtccacgta acgttgtctt    420
cgacatattt actggtaatg gacaagacat ggtgttcacc atttatggtg accattggcg    480
aaaaatgagg cgtattatga cgcttcatt tttcactaac aaagtgggtg accaatatag    540
tgatatgtgg gagaatgaga tggacttagt tgtaaatgac tgaagaaga atgaaaaagt    600
gaaatatgag ggaattgtga ttaggaaacg attgcagctg atgctgtata acatcatgta    660
tcgaatgatg tttgatgcca aatttgagtc ccaaatgat cctttgttca ttgaggcaac    720
aaagtttaat tcagagagaa gcagattagc tcagagcttt gactacaatt atggtgattt    780
tatcccttta cttagaccat tcttgagagg gtaccttaac aagtgtaaag acttacaac    840
aaggagactt gcattcttca acaattattt tgtagagaaa agaaggaaaa taatggatga    900
aaatggagaa aagcataaga taagctgtgc tattgatcac attatagatg ccgaaatgaa    960
aggagaaata aatgagcaaa atgtactcta tattgtggag aatatcaatg ttgcagcaat   1020
tgaaacaact ctatggtcca tggaatgggc catagctgaa cttgtaaatc atcccattgt   1080
tcaacagaag attagggatg aaatctcaac agtcctcaaa ggcagatcag tcacagaatc   1140
aaacctccat gagctgcctt acttgctagc aacagtaaat gaaacattaa gactccacac   1200
accaatacct ttacttgtac cccatgatga ccttgaagaa gcaaagttag gtggttacac   1260
tattcctaaa gaaactaagg tgggtgtgaa tgcgtggtgg ctggctaaca accctgcttg   1320
gtggaaaaac ccgaatgaat tccggcccga gaggtttctt gaggaggata gtagcacaga   1380
ggcagctggt gctggtggca aggtagattt caggtactta ccattcggta tggggaggcg   1440
gagctgcccc ggaatcatcc ttgcactgcc aattctgggg cttgtcatag ccaaactggt   1500
gtcaaatttt gaaatgcagg gtccaccagg tgtggaaaag gttgatacaa gtgaaagagg   1560
agggcagttt agcttgaca ttgcaaaaca ttccacggtt gtcttcaagc ctattgctgc   1620
ataataatat gcttaagcta tccttgtttt aattatattt gtcttaccag aaagcaaac   1680
tactaagtta ctcgataaag atttcaatga atattacagt ttttgttaaa aaaaaaaaaa   1740
aaaaa                                             1745
```

<210> SEQ ID NO 7
<211> LENGTH: 534
<212> TYPE: PRT
<213> ORGANISM: *Nicotiana tabacum*

<400> SEQUENCE: 7

Met 1	Ala	Lys	Leu	Leu 5	Asn	Asn	Thr	Ile	Phe 10	Cys	Ile	Leu	Phe	Ser 15	Ile
Val	Phe	Leu	Ser 20	Phe	Ala	Lys	Leu	Leu 25	Ser	Ser	Tyr	Leu	Ser 30	Ile	Pro
Phe	Pro	Leu 35	Glu	Tyr	Ile	Ser	Leu 40	Ile	Val	Leu	Leu 45	Leu	Pro	Leu	Ile
Ile	Asn 50	Phe	Leu	Cys	Val	Lys 55	Pro	Gln	Asn	Asn 60	Leu	Pro	Pro	Gly	Pro
Thr 65	Ala	Val	Pro	Ile	Phe 70	Gly	Asn	Trp	Leu	Gln 75	Val	Gly	Asn	Asp	Leu
Asn	His	Gln	Leu 85	Leu	Ala	Thr	Met	Ser 90	Gln	Thr	Tyr	Gly	Pro	Ile	Phe
Leu	Leu	Lys	Leu 100	Gly	Ser	Lys	Asn	Leu 105	Ala	Val	Val	Ser	Asn 110	Pro	Glu
Leu	Ala	Asn 115	Gln	Val	Leu	His	Thr 120	Gln	Gly	Val	Glu	Phe 125	Gly	Ser	Arg
Pro	Arg 130	Asn	Val	Val	Phe 135	Asp	Ile	Phe	Thr	Gly 140	Asn	Gly	Gln	Asp	Met
Val 145	Phe	Thr	Ile	Tyr 150	Gly	Asp	His	Trp	Arg	Lys 155	Met	Arg	Arg	Ile	Met
Thr	Leu	Pro	Phe 165	Phe	Thr	Asn	Lys	Val 170	Val	His	Gln	Tyr	Ser	Asp 175	Met
Trp	Glu	Asn	Glu 180	Met	Asp	Leu	Val 185	Val	Asp	Asp	Leu	Lys 190	Lys	Asn	Glu
Lys	Val	Lys 195	Tyr	Asp	Gly	Ile 200	Val	Ile	Arg	Lys	Arg	Leu 205	Gln	Leu	Met
Leu	Tyr 210	Asn	Ile	Met	Tyr 215	Arg	Met	Met	Phe	Asp 220	Ala	Lys	Phe	Glu	Ser
Gln 225	Asp	Asp	Pro	Leu	Phe 230	Ile	Glu	Ala	Thr	Lys 235	Phe	Asn	Ser	Glu	Arg
Ser	Arg	Leu	Ala 245	Gln	Ser	Phe	Asp	Tyr 250	Asn	Tyr	Gly	Asp	Phe	Ile 255	Pro
Leu	Leu	Arg	Pro 260	Phe	Leu	Lys	Gly 265	Tyr	Leu	Asn	Lys	Cys 270	Lys	Asp	Leu
Gln	Thr 275	Arg	Arg	Leu	Ala	Phe	Phe 280	Asn	Asn	Tyr	Phe	Val 285	Gly	Lys	Arg
Arg	Lys 290	Ile	Met	Gly	Glu	Asn 295	Gly	Glu	Lys	His 300	Lys	Ile	Cys	Cys	Ala
Ile 305	Asp	His	Ile	Ile	Asp 310	Ala	Glu	Met	Lys	Gly 315	Glu	Ile	Ser	Glu	Gln
Asn	Val	Leu	Tyr 325	Ile	Val	Glu	Asn	Ile 330	Asn	Val	Ala	Ala	Ile	Glu	Thr
Thr	Leu	Trp	Ser 340	Met	Glu	Trp	Ala 345	Ile	Ala	Glu	Leu	Val 350	Asn	His	Pro
Ile	Val 355	Gln	Lys	Ile	Arg	Asp 360	Glu	Ile	Ser	Thr	Val 365	Leu	Lys	Gly	
Lys	Ser 370	Val	Lys	Glu	Ser 375	Asn	Leu	His	Glu	Leu 380	Pro	Tyr	Leu	Leu	Ala
Thr 385	Val	Asn	Glu	Thr 390	Leu	Arg	Leu	His	Thr 395	Pro	Ile	Pro	Leu	Leu	Val
Pro	His	Met	Asn 405	Leu	Glu	Glu	Ala	Lys 410	Leu	Gly	Gly	Tyr	Thr	Ile 415	Pro
Lys	Glu	Thr 420	Lys	Val	Val	Val	Asn 425	Ala	Trp	Trp	Leu	Ala 430	Asn	Asn	Pro

-continued

Ala Trp Trp Lys Asn Gln Asn Glu Phe Arg Pro Glu Arg Phe Leu Glu
435 440 445

Glu Asp Ser Ser Thr Glu Ala Ala Val Ala Gly Gly Lys Val Asp Phe
450 455 460

Arg Tyr Leu Pro Phe Gly Met Gly Arg Arg Ser Cys Pro Gly Ile Ile
465 470 475 480

Leu Ala Leu Pro Ile Leu Gly Leu Val Ile Ala Lys Leu Val Ser Asn
485 490 495

Phe Glu Met Gln Ala Pro Pro Gly Val Gly Lys Val Asp Thr Ser Glu
500 505 510

Lys Gly Gly Gln Phe Ser Leu His Ile Ala Lys His Ser Thr Val Val
515 520 525

Phe Lys Pro Ile Ala Ala
530

<210> SEQ ID NO 8
<211> LENGTH: 1693
<212> TYPE: DNA
<213> ORGANISM: Nicotiana tabacum

<400> SEQUENCE: 8

cctctagcta attaaaaaca tggccaaact tctcaacaac accatctttt gcattctctt 60

ttcaattgta tttctttcat ttgccaaatt actatcctcc tacctctcta tacctttccc 120

tcttgagtac atttcactta ttgtcctttt acttccccta ataatcaact tcctctgtgt 180

taagccccaa aacaacctcc cacctgggcc aacagcagtc ccaatttttg gtaattggct 240

tcaagttggc aatgacttga accatcaact ccttgccacc atgtcacaaa cctatgggcc 300

tatatattta ctcaaaacttg gttcaaaaaa cctagctgtg gtatcgaacc ctgagctagc 360

taaccaagtt ctacacacgc aagggggtcg gtttgggtcc cgtccacgta acgttgtctt 420

tgatatattt actggtaatg gacaagacat ggtgttcacc atttatggtg accattggcg 480

aaaaatgagg cgtattatga cgcttcatt tttcactaac aaagtgggtg accaatatag 540

tgatatgtgg gagaatgaga tggacctagt tgttgatgac ttgaagaaga atgaaaaagt 600

gaaatatgac ggaattgtga ttaggaaacg attgcagctg atgctatata acattatgta 660

tcgaatgatg tttgatgcca agtttgagtc ccaagatgat cctttgttca ttgaggcaac 720

aaagtttaat tcagagagaa gcagattagc tcagagcttt gactacaatt atgggtgattt 780

tatccctttg cttagaccat tcttgaaagg gtaccttaac aagtgcacaa acttacaaac 840

aaggagactt gcattcttca acaattattt ttaggggaaa agaaggaaaa taatgggtga 900

aaatggagaa aaacacaaga tatgttgtgc tattgatcac attatagatg ctgaaatgaa 960

aggagaaata agtgagcaaa atgtactcta tattgtggag aatatcaatg ttgcagcaat 1020

tgaacaact ctatggtcca tggaaatggc catagctgag cttgtaaatc atcccattgt 1080

tcaacagaag attagggatg aaatctcaac agtcctcaaa ggaaagtcag tcaagaatc 1140

aaacctacat gagctgcctt acttgctagc aacagtaaat gaaacattaa gactccacac 1200

accaatacct ttacttgtac cacatatgaa ccttgaagaa gcaaagctag gtgggttacac 1260

tattcctaaa gaaactaagg tgggtgtgaa tgcattgggt ctggctaaca accctgcctg 1320

gtggaaaaac cagaacgaat tccggcccg gcggtttctc gaggaggata gtagcacaga 1380

ggcagctggt gctggtggca aggttgattt caggtacttg cccttcggtg tggggaggcg 1440

gagctgcccc ggaatcatcc ttgcactgcc aattctgggg cttgtcatag ccaaaactgt 1500

-continued

```

gtcaaatttt gaaatgcagg ctccctccagg tgtaggaaaa gttgatacaa gtgagaaagg 1560
agggcagttt agcttgcaca ttgcaaaaca ttccacggtt gtcttcaagc ctattgctgc 1620
ataatattac agtttttgtt actctataaa gatttcaatg aatattacag tttttgttaa 1680
aaaaaaaaa aaa 1693

```

```

<210> SEQ ID NO 9
<211> LENGTH: 519
<212> TYPE: PRT
<213> ORGANISM: Nicotiana tabacum

```

```

<400> SEQUENCE: 9

```

```

Met Tyr His Leu Leu Ser Pro Ile Glu Ala Ile Val Gly Leu Val Thr
1           5           10           15
Phe Ala Phe Leu Leu Tyr Leu Leu Trp Thr Lys Lys Gln Ser Lys Ile
20           25           30
Leu Asn Pro Leu Pro Pro Lys Ile Pro Gly Gly Trp Pro Val Ile Gly
35           40           45
His Leu Phe Tyr Phe Asn Asn Asn Gly Asp Asp Asp Arg His Phe Ser
50           55           60
Gln Lys Leu Gly Asp Leu Ala Asp Lys Tyr Gly Pro Val Phe Thr Phe
65           70           75           80
Arg Leu Gly Phe Arg Arg Phe Leu Ala Val Ser Ser Tyr Glu Ala Met
85           90           95
Lys Glu Cys Phe Ser Thr Asn Asp Ile His Phe Ala Asp Arg Pro Ala
100          105          110
Leu Leu Tyr Gly Glu Tyr Leu Cys Tyr Asn Asn Ala Met Leu Ala Val
115          120          125
Ala Lys Tyr Gly Pro Tyr Trp Lys Lys Asn Arg Lys Leu Val Asn Gln
130          135          140
Glu Leu Leu Ser Val Ser Arg Leu Glu Lys Phe Lys His Val Arg Phe
145          150          155          160
Ser Ile Val Gln Lys Asn Ile Lys Gln Leu Tyr Asn Cys Asp Ser Pro
165          170          175
Met Val Lys Ile Asn Leu Ser Asp Trp Ile Asp Lys Leu Thr Phe Asp
180          185          190
Ile Ile Leu Lys Met Val Val Gly Lys Thr Tyr Asn Asn Gly His Gly
195          200          205
Glu Ile Leu Lys Ala Ala Phe Gln Lys Phe Met Val Gln Ala Met Glu
210          215          220
Ile Glu Leu Tyr Asp Val Phe His Ile Pro Phe Phe Lys Trp Leu Asp
225          230          235          240
Leu Thr Gly Asn Ile Lys Ala Met Lys Gln Thr Phe Lys Asp Ile Asp
245          250          255
Asn Ile Ile Gln Gly Trp Leu Asp Glu His Ile Lys Lys Arg Glu Thr
260          265          270
Lys Asp Val Gly Gly Glu Asn Glu Gln Asp Phe Ile Asp Val Leu Leu
275          280          285
Ser Lys Arg Ser Asn Glu His Leu Gly Asp Gly Tyr Ser His Asp Thr
290          295          300
Thr Ile Lys Ala Thr Val Phe Thr Leu Val Leu Asp Ala Thr Asp Thr
305          310          315          320
Leu Ala Leu His Ile Lys Trp Val Met Ala Leu Met Ile Asn Asn Lys
325          330          335
Asn Val Met Lys Lys Ala Gln Glu Glu Met Asp Thr Ile Val Gly Arg

```


-continued

340	345	350
Asp Arg Trp Val Glu Glu Asn Asp 355	Ile Lys Asn Leu Val Tyr Leu Gln 360	
Ala Ile Val Lys Glu Val Leu Arg Leu His Pro 370	Pro Ala Pro Leu Ser 380	
Val Gln His Leu Ser Val Lys Asp Cys Val 385	Val Asn Gly Tyr His Ile 395	400
Pro Lys Gly Thr Ala Leu Leu Thr Asn Ile Met 405	Lys Leu Gln Arg Asp 410	415
Pro Gln Ile Trp Val Asp Pro Asp Thr Phe Asp 420	Pro Glu Arg Phe Leu 425	430
Thr Thr Asn Ala Ala Ile Asp Tyr Arg Gly Gln His 435	Tyr Glu Leu Ile 440	445
Pro Phe Gly Ser Gly Arg Arg Ala Cys Pro Ala 450	Met Asn Tyr Ser Leu 455	460
Gln Val Glu His Leu Ser Ile Ala His Leu Ile 465	Gln Gly Phe Asn Phe 470	475
Ala Thr Thr Thr Asn Glu Pro Leu Asp Met Lys 485	Gln Gly Val Gly Leu 490	495
Thr Leu Pro Lys Lys Thr Asp Val Glu Val Leu Ile 500	Thr Pro Arg Leu 505	510
Pro Pro Thr Leu Tyr Gln Tyr 515		

<210> SEQ ID NO 10

<211> LENGTH: 1578

<212> TYPE: DNA

<213> ORGANISM: Nicotiana tabacum

<400> SEQUENCE: 10

```

atgtatcatc ttctttctcc catagaagcc attgtaggac ttgtaacctt tgcattttcta    60
ctctacttgc tatggacaaa aaaacaatca aaaatcttaa acccactgcc tccaaaaatc    120
ccaggtggat ggccagtaat cggccatctc ttttatttca acaacaatgg cgatgatgac    180
cgccattttt ctcaaaaact cgggagactta gctgacaaat atgggtcccggt cttcacattc    240
cggttagggg ttgcgcgttt cttggcgggt agtagttatg aagctatgaa agaatgcttc    300
tctaccaatg atatccattt cgccgatcgg ccagctttac tttacggaga atacctttgc    360
tataacaatg ccattgcttc tgttgccaaa tatggccctt actggaaaaa aaatcgaaag    420
ctagtcaatc aagaacttct ctccgttagt cggctcgaaa aattcaaaca tgttagattt    480
tctatagttc agaaaaatat taaacaattg tataattgtg attcaccaat ggtgaagata    540
aaccttagtg attgtagata taaattgaca ttcgacatca ttttgaatat ggttgttggg    600
aagacctata ataattggaca tggagaaata ctcaaagcag cttttcagaa gttcatgggt    660
caagctatgg agattgagct ctatgatggt tttcacattc catttttcaa gtggttggtg    720
cttacaggga atattaaggc tatgaaacaa actttcaaag acattgataa tattatccaa    780
ggttgggttag atgagacacat taagaagaga gaaacaaagg atgttgaggg tgaataatgaa    840
caagatttta ttgatgtgct gctttccaag aggagcaacg aacatcttgg cgatgggttac    900
tctcatgaca ccaccatcaa agcaacagta ttcactttgg tcttggtatgc aacagacaca    960
cttgcacttc atataaagtg ggtaattggc ttaatgataa acaataagaa tgtcatgaag    1020
aaagcacaag aagagatgga caccattggt ggtagagata gatgggtaga agagaatgat    1080
atcaagaatt tgggtgtatct tcaagcaatt gttaaagaag tattacgatt acatccacct    1140

```

-continued

```

gcacctttgt cagtacaaca cctatccgta aaagattgtg ttgtcaatgg ataccatatt 1200
cctaagggga ctgcactact tacaaatatt atgaaacttc aacgagaccc acaaatatgg 1260
gtagatcctg atacattcga tccagaaaga ttcttgacga ctaatgctgc aattgactat 1320
cgcgggcagc actatgagtt gatcccgttt ggatcagggg gacgagcttg tcccgcgatg 1380
aattactcat tgcaagtgga acacctttca attgctcatt tgatccaggg tttcaatttt 1440
gcaactacga ctaacgagcc tttggatatg aaacaaggcg tgggtctaac tttacctaag 1500
aagacagatg ttgaagtgct aattacacct cgccttcctc ctacgcttta tcaatattaa 1560
tatgttttgt tgttgtga 1578

```

```

<210> SEQ ID NO 11
<211> LENGTH: 509
<212> TYPE: PRT
<213> ORGANISM: Nicotiana tabacum

```

```

<400> SEQUENCE: 11

```

```

Met Glu Gly Thr Asn Leu Thr Thr Tyr Ala Ala Val Phe Leu Gly Thr
1           5           10           15
Leu Phe Leu Leu Leu Leu Ser Lys Phe Leu Arg Gln Arg Lys Leu Asn
20          25          30
Leu Pro Pro Gly Pro Lys Pro Trp Pro Ile Ile Gly Asn Leu Asn Leu
35          40          45
Ile Gly Asn Leu Pro His Arg Ser Ile His Glu Leu Ser Leu Lys Tyr
50          55          60
Gly Pro Ile Met Gln Leu Gln Phe Gly Thr Phe Pro Val Val Val Gly
65          70          75          80
Ser Ser Val Glu Met Ala Lys Val Phe Leu Lys Ser Met Asp Ile Asn
85          90          95
Phe Val Gly Arg Pro Lys Thr Ala Ala Gly Lys Tyr Thr Thr Tyr Asn
100         105         110
Tyr Ser Asp Ile Thr Trp Ser Pro Tyr Gly Pro Tyr Trp Arg Gln Ala
115         120         125
Arg Arg Met Cys Leu Met Glu Leu Phe Ser Thr Lys Arg Leu Asp Ser
130         135         140
Tyr Glu Tyr Ile Arg Ala Glu Glu Leu His Ser Leu Leu His Asn Leu
145         150         155         160
Asn Lys Ile Ser Gly Lys Pro Ile Val Leu Lys Asp Tyr Leu Thr Thr
165         170         175
Leu Ser Leu Asn Val Ile Ser Arg Met Val Leu Gly Lys Arg Tyr Leu
180         185         190
Asp Glu Ser Glu Asn Ser Ile Val Thr Pro Glu Glu Phe Lys Lys Met
195         200         205
Leu Asp Glu Leu Phe Leu Leu Asn Gly Val Leu Asn Ile Gly Asp Ser
210         215         220
Ile Pro Trp Ile Asp Phe Met Asp Leu Gln Gly Tyr Val Lys Arg Met
225         230         235         240
Lys Phe Val Ser Lys Lys Phe Asp Lys Phe Leu Glu His Val Ile Asp
245         250         255
Glu His Asn Val Arg Arg Asn Gly Val Glu Asn Tyr Ile Ala Lys Asp
260         265         270
Met Val Asp Val Leu Leu Gln Leu Ala Asp Asp Pro Thr Leu Glu Val
275         280         285
Lys Leu Glu Arg His Gly Val Lys Ala Phe Thr Gln Asp Met Leu Ala

```

-continued

290	295	300
Gly Gly Thr Glu Ser Ser Ala Val Thr Val Glu Trp Ala Ile Ser Glu 305 310 315 320		
Leu Leu Lys Lys Pro Glu Ile Phe Lys Lys Ala Thr Glu Glu Leu Asp 325 330 335		
Arg Val Ile Gly Gln Asn Arg Trp Val Gln Glu Lys Asp Ile Pro Asn 340 345 350		
Leu Pro Tyr Ile Glu Ala Ile Val Lys Glu Thr Met Arg Leu His Pro 355 360 365		
Val Ala Pro Met Leu Val Pro Arg Glu Cys Arg Glu Asp Cys Lys Val 370 375 380		
Ala Gly Tyr Asp Val Lys Lys Gly Thr Arg Val Leu Val Ser Val Trp 385 390 395 400		
Thr Ile Gly Arg Asp Pro Thr Leu Trp Asp Glu Pro Glu Ala Phe Lys 405 410 415		
Pro Glu Arg Phe His Glu Lys Ser Ile Asp Val Lys Gly His Asp Phe 420 425 430		
Glu Leu Leu Pro Phe Gly Ala Gly Arg Arg Met Cys Pro Gly Tyr Asn 435 440 445		
Leu Gly Leu Lys Val Ile Gln Ala Ser Leu Ala Asn Leu Ile His Gly 450 455 460		
Phe Asn Trp Ser Leu Pro Asp Asn Met Thr Pro Glu Asp Leu Asp Met 465 470 475 480		
Asp Glu Ile Phe Gly Leu Ser Thr Pro Lys Lys Phe Pro Leu Ala Thr 485 490 495		
Val Ile Glu Pro Arg Leu Ser Pro Lys Leu Tyr Ser Val 500 505		

<210> SEQ ID NO 12

<211> LENGTH: 1530

<212> TYPE: DNA

<213> ORGANISM: Nicotiana tabacum

<400> SEQUENCE: 12

atggaaggta caaacttgac tacatatgca gcagtatttc ttggtactct gtttcttttg	60
ctcctttcca aatttcttcg ccaaagaaaa ctcaacttac ctccaggccc aaaaccatgg	120
ccgatcatcg gaaacttaaa cottatcggc aatcttcttc atcgctcaat ccacgaactt	180
tcaactcaagt acgggccaat tatgcaactc caattcggga ctttccccgt tgtgtgtggc	240
tcttcgctcg aaatggccaa ggttttcttc aaatcaatgg atattaactt tgtaggcagg	300
cctaaaacgg ccgcccggaa gtacacaact tacaattatt cagatattac atggtctcct	360
tatggaccat attggcgcca ggcacgtaga atgtgcctaa tggaattatt cagcacgaaa	420
cgtctcgatt catacagtaga tattcgggct gaggagtgc attctctgct ccataatttg	480
aataaaaatat cagggaaacc aattgtgctg aaagattatt tgacgacgtt gagtttaa	540
gttattagca ggatgttact ggggaagagg tatttggacg aatccgagaa ctcgattgtg	600
actcctgagg aatttaagaa gatgttgac gagctgttct tgctaaatgg tgtactta	660
attggagatt caattccctg gattgatttc atggacttac aaggttatgt taagaggatg	720
aaatttgtga gcaagaaatt cgacaagttt ttggagcatg ttatcgatga gcataacgtt	780
aggagaaatg gattggagaa ttacattgct aaggacatgg ttgatgttct gttgcagctt	840
gctgatgac cgacgttgga agttaagctg gagagacatg gactcaaagc attcactcag	900
gatatgcttg ctggtggaac cgagagttca gcagtgcagc tggagtgggc aatttcggag	960

-continued

```

ctgctaaaga agccagagat tttcaaaaag gctacagaag aattggatcg agtaattggg 1020
cagaatagat gggtagaaga aaaagacatt ccaaatcttc cttacataga ggcaatagtc 1080
aaagagacta tgcgactgca ccccgtagga ccaatgttgg tgccacggga gtgtcgagaa 1140
gactgtaagg tagcaggcta cgacgttaag aaaggaacca gggtccttgt gagegtatgg 1200
actattggaa gagaccctac attgtgggac gagcctgagg cgttcaagcc ggagaggttc 1260
cacgaaaagt ccattgatgt taaaggacat gattttgagc ttttgccatt tggagctggg 1320
agaaggatgt gcccgggtta taacttgggg cttaagggtga ttcaagctag cttagctaatt 1380
cttatacatg gatttaactg gtcattgcct gataatatga ctctgagga cctcgacatg 1440
gatgagatgt ttgggtcttc cacacctaaa aagtttcac ttgctactgt gattgagcca 1500
agactttcac caaaacttta ctctgtttga 1530

```

```

<210> SEQ ID NO 13
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Derived from Nicotiana tabacum p450 gene

```

```

<400> SEQUENCE: 13

```

```

ggcggagaat ttgtcctgga atgtcatttg gtttag 36

```

```

<210> SEQ ID NO 14
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Derived from Nicotiana tabacum p450 gene

```

```

<400> SEQUENCE: 14

```

```

gtacaatagt gaggttgaca atg 23

```

```

<210> SEQ ID NO 15
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Derived from Nicotiana tabacum p450 gene

```

```

<400> SEQUENCE: 15

```

```

ggtggttgat aatgcatg 18

```

```

<210> SEQ ID NO 16
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Derived from Nicotiana tabacum p450 gene

```

```

<400> SEQUENCE: 16

```

```

ttatgcagca ataggcttga agaca 25

```

```

<210> SEQ ID NO 17
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Derived from Nicotiana tabacum p450 gene

```

```

<400> SEQUENCE: 17

```

```

gggggatcca tgcaattctt cagcttggtt tcc 33

```

-continued

<210> SEQ ID NO 18
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Derived from Nicotiana tabacum p450 gene

<400> SEQUENCE: 18

ggggaattct tactctcgag aaggttgata agg 33

<210> SEQ ID NO 19
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Derived from Nicotiana tabacum p450 gene

<400> SEQUENCE: 19

cccggatcca tgtatcatct tctttctccc 30

<210> SEQ ID NO 20
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Derived from Nicotiana tabacum p450 gene

<400> SEQUENCE: 20

ggggaattct caatattgat aaagcgtagg agg 33

<210> SEQ ID NO 21
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Derived from Nicotiana tabacum p450 gene

<400> SEQUENCE: 21

cccggatcca tgcaatcctt cagcttggtt tcc 33

<210> SEQ ID NO 22
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Derived from Nicotiana tabacum p450 gene

<400> SEQUENCE: 22

ggggagctct cactcgcaag aagattgata agg 33

<210> SEQ ID NO 23
<211> LENGTH: 75
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Derived from Nicotiana tabacum p450 gene

<400> SEQUENCE: 23

gccattatcg gcgcaatact aatctccaaa ctccgcggtg aaaaattcaa gctcccacct 60

ggtccaacag cagtc 75

<210> SEQ ID NO 24
<211> LENGTH: 75
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Derived from Nicotiana tabacum p450 gene

<400> SEQUENCE: 24

gggggatcca tggacctcct cctcatagaa aaaaccctcg tcgccttatt cgcgcgcatt 60
atcggcgcaa tacta 75

<210> SEQ ID NO 25
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Derived from Nicotiana tabacum p450 gene

<400> SEQUENCE: 25

ggggagctct tatgcagcaa taggcttgaa gac 33

<210> SEQ ID NO 26
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Nicotiana tabacum
<220> FEATURE:
<223> OTHER INFORMATION: Nicotiana tabacum p450 protein
<220> FEATURE:
<221> NAME/KEY: Variant
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa = any amino acid

<400> SEQUENCE: 26

Lys Glu Thr Leu Arg Leu Xaa
1 5

<210> SEQ ID NO 27
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Nicotiana tabacum
<220> FEATURE:
<223> OTHER INFORMATION: Nicotiana tabacum p450 protein
<220> FEATURE:
<221> NAME/KEY: Variant
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa = any amino acid
<220> FEATURE:
<221> NAME/KEY: Variant
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa = any amino acid

<400> SEQUENCE: 27

Pro Phe Gly Xaa Gly Arg Arg Xaa Cys Pro Ala
1 5 10

<210> SEQ ID NO 28
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Nicotiana tabacum
<220> FEATURE:
<223> OTHER INFORMATION: Nicotiana tabacum p450 protein
<220> FEATURE:
<221> NAME/KEY: Variant
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa = any amino acid
<220> FEATURE:
<221> NAME/KEY: Variant
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa = any amino acid

<400> SEQUENCE: 28

Pro Phe Gly Xaa Gly Arg Arg Xaa Cys Pro Gly
1 5 10

-continued

<210> SEQ ID NO 29
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Nicotiana tabacum
<220> FEATURE:
<223> OTHER INFORMATION: Nicotiana tabacum p450 protein
<220> FEATURE:
<221> NAME/KEY: Variant
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa = any amino acid

<400> SEQUENCE: 29

Phe Xaa Pro Glu Arg Phe
1 5

<210> SEQ ID NO 30
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Derived from Nicotiana tabacum p450 protein

<400> SEQUENCE: 30

Ala Ala Arg Gly Ala Arg Ala Cys Ile Tyr Thr Ile Met Gly Ile Tyr
1 5 10 15

Thr Ile Cys Ala
20

<210> SEQ ID NO 31
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Derived from Nicotiana tabacum p450 protein

<400> SEQUENCE: 31

Ala Ala Arg Gly Ala Arg Ala Cys Ile Tyr Thr Ile Met Gly Ile Tyr
1 5 10 15

Thr Ile Thr Ala
20

<210> SEQ ID NO 32
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Derived from Nicotiana tabacum p450 protein

<400> SEQUENCE: 32

Ala Ala Arg Gly Ala Arg Ala Cys Ile Tyr Thr Ile Met Gly Ile Tyr
1 5 10 15

Thr Ile Met Gly
20

<210> SEQ ID NO 33
<211> LENGTH: 18
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Derived from Nicotiana tabacum p450 protein

<400> SEQUENCE: 33

Thr Thr Tyr Ile Ile Ile Cys Cys Ile Gly Ala Arg Met Gly Ile Thr
1 5 10 15

Thr Tyr

-continued

<210> SEQ ID NO 34
<211> LENGTH: 18
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Derived from Nicotiana tabacum p450 protein

<400> SEQUENCE: 34

Arg Ala Ala Ile Cys Lys Tyr Thr Cys Ile Gly Gly Ile Ile Ile Arg
1 5 10 15

Ala Ala

<210> SEQ ID NO 35
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Derived from Nicotiana tabacum p450 protein

<400> SEQUENCE: 35

Gly Gly Ile Met Gly Ile Met Gly Ile Ile Ile Ile Thr Gly Tyr Cys
1 5 10 15

Cys Ile Gly Ser
20

<210> SEQ ID NO 36
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Derived from Nicotiana tabacum p450 protein

<400> SEQUENCE: 36

Cys Lys Ile Cys Lys Ile Cys Cys Ile Ile Ile Ile Cys Cys Arg Ala
1 5 10 15

Ala Ile Gly Gly
20

<210> SEQ ID NO 37
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Derived from T7 bacteriophage promoter

<400> SEQUENCE: 37

gtaatacgcac tcactatagg g

21

<210> SEQ ID NO 38
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Derived from T3 bacteriophage promoter

<400> SEQUENCE: 38

caattaaccc tcactaaagg g

21

<210> SEQ ID NO 39
<211> LENGTH: 500
<212> TYPE: PRT
<213> ORGANISM: Mentha piperita

<400> SEQUENCE: 39

Met Glu Leu Gln Ile Ser Ser Ala Ile Ile Ile Leu Val Val Thr Tyr

-continued

1	5	10	15
Thr Ile Ser Leu Leu Ile Ile Lys Gln Trp Arg Lys Pro Lys Pro Gln	20	25	30
Glu Asn Leu Pro Pro Gly Pro Pro Lys Leu Pro Leu Ile Gly His Leu	35	40	45
His Leu Leu Trp Gly Lys Leu Pro Gln His Ala Leu Ala Ser Val Ala	50	55	60
Lys Gln Tyr Gly Pro Val Ala His Val Gln Leu Gly Glu Val Phe Ser	65	70	75
Val Val Leu Ser Ser Arg Glu Ala Thr Lys Glu Ala Met Lys Leu Val	85	90	95
Asp Pro Ala Cys Ala Asp Arg Phe Glu Ser Ile Gly Thr Lys Ile Met	100	105	110
Trp Tyr Asp Asn Asp Asp Ile Ile Phe Ser Pro Tyr Ser Val His Trp	115	120	125
Arg Gln Met Arg Lys Ile Cys Val Ser Glu Leu Leu Ser Ala Arg Asn	130	135	140
Val Arg Ser Phe Gly Phe Ile Arg Gln Asp Glu Val Ser Arg Leu Leu	145	150	155
Gly His Leu Arg Ser Ser Ala Ala Ala Gly Glu Ala Val Asp Leu Thr	165	170	175
Glu Arg Ile Ala Thr Leu Thr Cys Ser Ile Ile Cys Arg Ala Ala Phe	180	185	190
Gly Ser Val Ile Arg Asp His Glu Glu Leu Val Glu Leu Val Lys Asp	195	200	205
Ala Leu Ser Met Ala Ser Gly Phe Glu Leu Ala Asp Met Phe Pro Ser	210	215	220
Ser Lys Leu Leu Asn Leu Leu Cys Trp Asn Lys Ser Lys Leu Trp Arg	225	230	235
Met Arg Arg Arg Val Asp Ala Ile Leu Glu Ala Ile Val Glu Glu His	245	250	255
Lys Leu Lys Lys Ser Gly Glu Phe Gly Gly Glu Asp Ile Ile Asp Val	260	265	270
Leu Phe Arg Met Gln Lys Asp Ser Gln Ile Lys Val Pro Ile Thr Thr	275	280	285
Asn Ala Ile Lys Ala Phe Ile Phe Asp Thr Phe Ser Ala Gly Thr Glu	290	295	300
Thr Ser Ser Thr Thr Thr Leu Trp Val Met Ala Glu Leu Met Arg Asn	305	310	315
Pro Glu Val Met Ala Lys Ala Gln Ala Glu Val Arg Ala Ala Leu Lys	325	330	335
Gly Lys Thr Asp Trp Asp Val Asp Asp Val Gln Glu Leu Lys Tyr Met	340	345	350
Lys Ser Val Val Lys Glu Thr Met Arg Met His Pro Pro Ile Pro Leu	355	360	365
Ile Pro Arg Ser Cys Arg Glu Glu Cys Glu Val Asn Gly Tyr Thr Ile	370	375	380
Pro Asn Lys Ala Arg Ile Met Ile Asn Val Trp Ser Met Gly Arg Asn	385	390	395
Pro Leu Tyr Trp Glu Lys Pro Glu Thr Phe Trp Pro Glu Arg Phe Asp	405	410	415
Gln Val Ser Arg Asp Phe Met Gly Asn Asp Phe Glu Phe Ile Pro Phe	420	425	430

[illegible]

```
<210> SEQ ID NO 40
<211> LENGTH: 496
<212> TYPE: PRT
<213> ORGANISM: Mentha spicata
```

<400> SEQUENCE: 40

Met 1	Glu	Leu	Asp	Leu 5	Leu	Ser	Ala	Ile	Ile 10	Ile	Leu	Val	Ala	Thr 15	Tyr
Ile	Val	Ser	Leu 20	Leu	Ile	Asn	Gln	Trp 25	Arg	Lys	Ser	Lys	Ser 30	Gln	Gln
Asn	Leu	Pro 35	Pro	Ser	Pro	Pro	Lys 40	Leu	Pro	Val	Ile 45	Gly	His	Leu	His
Phe 50	Leu	Trp	Gly	Gly	Leu	Pro 55	Gln	His	Val	Phe	Arg 60	Ser	Ile	Ala	Gln
Lys 65	Tyr	Gly	Pro	Val	Ala 70	His	Val	Gln	Leu	Gly 75	Glu	Val	Tyr	Ser	Val 80
Val	Leu	Ser	Ser 85	Ala	Glu	Ala	Ala	Lys 90	Gln	Ala	Met	Lys	Val	Leu 95	Asp
Pro	Asn	Phe	Ala 100	Asp	Arg	Phe	Asp	Gly 105	Ile	Gly	Ser	Arg	Thr 110	Met	Trp
Tyr	Asp	Lys 115	Asp	Asp	Ile	Ile	Phe 120	Ser	Pro	Tyr	Asn	Asp 125	His	Trp	Arg
Gln 130	Met	Arg	Arg	Ile	Cys 135	Val	Thr	Glu	Leu	Leu	Ser 140	Pro	Lys	Asn	Val
Arg 145	Ser	Phe	Gly	Tyr	Ile 150	Arg	Gln	Glu	Glu	Ile 155	Glu	Arg	Leu	Ile	Arg 160
Leu	Leu	Gly	Ser 165	Ser	Gly	Gly	Ala	Pro	Val	Asp 170	Val	Thr	Glu	Glu 175	Val
Ser	Lys	Met 180	Ser	Cys	Val	Val	Val	Cys 185	Arg	Ala	Ala	Phe	Gly 190	Ser	Val
Leu	Lys	Asp 195	Gln	Gly	Ser	Leu	Ala 200	Glu	Leu	Val	Lys	Glu 205	Ser	Leu	Ala
Leu 210	Ala	Ser	Gly	Phe	Glu 215	Leu	Ala	Asp	Leu	Tyr 220	Pro	Ser	Ser	Trp	Leu
Leu 225	Asn	Leu	Leu	Ser	Leu 230	Asn	Lys	Tyr	Arg	Leu 235	Gln	Arg	Met	Arg	Arg 240
Arg	Leu	Asp	His 245	Ile	Leu	Asp	Gly	Phe	Leu	Glu 250	Glu	His	Arg	Glu 255	Lys
Lys	Ser	Gly 260	Glu	Phe	Gly	Gly	Glu 265	Asp	Ile	Val	Asp	Val	Leu 270	Phe	Arg
Met	Gln	Lys 275	Gly	Ser	Asp	Ile	Lys 280	Ile	Pro	Ile	Thr	Ser 285	Asn	Cys	Ile
Lys 290	Gly	Phe	Ile	Phe	Asp 295	Thr	Phe	Ser	Ala	Gly 300	Ala	Glu	Thr	Ser	Ser

-continued

Thr Thr Ile Ser Trp Ala Leu Ser Glu Leu Met Arg Asn Pro Ala Lys
 305 310 315 320
 Met Ala Lys Val Gln Ala Glu Val Arg Glu Ala Leu Lys Gly Lys Thr
 325 330 335
 Val Val Asp Leu Ser Glu Val Gln Glu Leu Lys Tyr Leu Arg Ser Val
 340 345 350
 Leu Lys Glu Thr Leu Arg Leu His Pro Pro Phe Pro Leu Ile Pro Arg
 355 360 365
 Gln Ser Arg Glu Glu Cys Glu Val Asn Gly Tyr Thr Ile Pro Ala Lys
 370 375 380
 Thr Arg Ile Phe Ile Asn Val Trp Ala Ile Gly Arg Asp Pro Gln Tyr
 385 390 395 400
 Trp Glu Asp Pro Asp Thr Phe Arg Pro Glu Arg Phe Asp Glu Val Ser
 405 410 415
 Arg Asp Phe Met Gly Asn Asp Phe Glu Phe Ile Pro Phe Gly Ala Gly
 420 425 430
 Arg Arg Ile Cys Pro Gly Leu His Phe Gly Leu Ala Asn Val Glu Ile
 435 440 445
 Pro Leu Ala Gln Leu Leu Tyr His Phe Asp Trp Lys Leu Pro Gln Gly
 450 455 460
 Met Thr Asp Ala Asp Leu Asp Met Thr Glu Thr Pro Gly Leu Ser Gly
 465 470 475 480
 Pro Lys Lys Lys Asn Val Cys Leu Val Pro Thr Leu Tyr Lys Ser Pro
 485 490 495

<210> SEQ ID NO 41

<211> LENGTH: 509

<212> TYPE: PRT

<213> ORGANISM: *Nepeta racemosa*

<400> SEQUENCE: 41

Met Val Ser Leu Ser Tyr Phe Leu Ile Ala Leu Leu Cys Thr Leu Pro
 1 5 10 15
 Phe Leu Leu Phe Leu Asn Lys Trp Arg Arg Ser Tyr Ser Gly Lys Thr
 20 25 30
 Pro Pro Pro Ser Pro Pro Lys Leu Pro Val Ile Gly Asn Leu His Gln
 35 40 45
 Leu Gly Leu Tyr Pro His Arg Tyr Leu Gln Ser Leu Ser Arg Arg Tyr
 50 55 60
 Gly Pro Leu Met Gln Leu His Phe Gly Ser Val Pro Val Leu Val Ala
 65 70 75 80
 Ser Ser Pro Glu Ala Ala Arg Glu Ile Met Lys Asn Gln Asp Ile Val
 85 90 95
 Phe Ser Asn Arg Pro Lys Met Ser Ile Ala Asn Arg Leu Phe Phe Asn
 100 105 110
 Asn Arg Asp Val Ala Phe Thr Gln Tyr Gly Glu Tyr Trp Arg Gln Ile
 115 120 125
 Arg Ser Ile Cys Val Leu Gln Leu Leu Ser Asn Lys Arg Val Gln Ser
 130 135 140
 Phe Arg Arg Val Arg Glu Glu Glu Thr Ser Ile Met Val Glu Lys Ile
 145 150 155 160
 Met Gln Leu Gly Ser Ser Ser Ser Thr Pro Val Asn Leu Ser Glu Leu
 165 170 175
 Leu Leu Ser Leu Thr Asn Asp Val Val Cys Arg Val Thr Leu Gly Lys
 180 185 190

-continued

Lys Tyr Gly Gly Gly Asn Gly Ser Glu Glu Val Asp Lys Leu Lys Glu
 195 200 205
 Met Leu Thr Glu Ile Gln Asn Leu Met Gly Ile Ser Pro Val Trp Glu
 210 215 220
 Phe Ile Pro Trp Leu Asn Trp Thr Arg Arg Phe Asp Gly Val Asp Gln
 225 230 235 240
 Arg Val Asp Arg Ile Val Lys Ala Phe Asp Gly Phe Leu Glu Ser Val
 245 250 255
 Ile Gln Glu His Lys Glu Arg Asp Gly Asp Lys Asp Gly Asp Gly Asp
 260 265 270
 Gly Ala Leu Asp Phe Val Asp Ile Leu Leu Gln Phe Gln Arg Glu Asn
 275 280 285
 Lys Asn Arg Ser Pro Val Glu Asp Asp Thr Val Lys Ala Leu Ile Leu
 290 295 300
 Asp Met Phe Val Ala Gly Thr Asp Thr Thr Ala Thr Ala Leu Glu Trp
 305 310 315 320
 Ala Val Ala Glu Leu Ile Lys Asn Pro Arg Ala Met Lys Arg Leu Gln
 325 330 335
 Asn Glu Val Arg Glu Val Ala Gly Ser Lys Ala Glu Ile Glu Glu Glu
 340 345 350
 Asp Leu Glu Lys Met Pro Tyr Leu Lys Ala Ser Ile Lys Glu Ser Leu
 355 360 365
 Arg Leu His Val Pro Val Val Leu Leu Val Pro Arg Glu Ser Thr Arg
 370 375 380
 Asp Thr Asn Val Leu Gly Tyr Asp Ile Ala Ser Gly Thr Arg Val Leu
 385 390 395 400
 Ile Asn Ala Trp Ala Ile Ala Arg Asp Pro Ser Val Trp Glu Asn Pro
 405 410 415
 Glu Glu Phe Leu Pro Glu Arg Phe Leu Asp Ser Ser Ile Asp Tyr Lys
 420 425 430
 Gly Leu His Phe Glu Leu Leu Pro Phe Gly Ala Gly Arg Arg Gly Cys
 435 440 445
 Pro Gly Ala Thr Phe Ala Val Ala Ile Asp Glu Leu Ala Leu Ala Lys
 450 455 460
 Leu Val His Lys Phe Asp Phe Gly Leu Pro Asn Gly Ala Arg Met Glu
 465 470 475 480
 Glu Leu Asp Met Ser Glu Thr Ser Gly Met Thr Val His Lys Lys Ser
 485 490 495
 Pro Leu Leu Leu Leu Pro Ile Pro His His Ala Ala Pro
 500 505

<210> SEQ ID NO 42

<211> LENGTH: 471

<212> TYPE: PRT

<213> ORGANISM: *Persea americana*

<400> SEQUENCE: 42

Met Ala Ile Leu Val Ser Leu Leu Phe Leu Ala Ile Ala Leu Thr Phe
 1 5 10 15
 Phe Leu Leu Lys Leu Asn Glu Lys Arg Glu Lys Lys Pro Asn Leu Pro
 20 25 30
 Pro Ser Pro Pro Asn Leu Pro Ile Ile Gly Asn Leu His Gln Leu Gly
 35 40 45
 Asn Leu Pro His Arg Ser Leu Arg Ser Leu Ala Asn Glu Leu Gly Pro
 50 55 60

-continued

Leu	Ile	Leu	Leu	His	Leu	Gly	His	Ile	Pro	Thr	Leu	Ile	Val	Ser	Thr	65	70	75	80
Ala	Glu	Ile	Ala	Glu	Glu	Ile	Leu	Lys	Thr	His	Asp	Leu	Ile	Phe	Ala	85	90	95	
Ser	Arg	Pro	Ser	Thr	Thr	Ala	Ala	Arg	Arg	Ile	Phe	Tyr	Asp	Cys	Thr	100	105	110	
Asp	Val	Ala	Phe	Ser	Pro	Tyr	Gly	Glu	Tyr	Trp	Arg	Gln	Val	Arg	Lys	115	120	125	
Ile	Cys	Val	Leu	Glu	Leu	Leu	Ser	Ile	Lys	Arg	Val	Asn	Ser	Tyr	Arg	130	135	140	
Ser	Ile	Arg	Glu	Glu	Glu	Val	Gly	Leu	Met	Met	Glu	Arg	Ile	Ser	Gln	145	150	155	160
Ser	Cys	Ser	Thr	Gly	Glu	Ala	Val	Asn	Leu	Ser	Glu	Leu	Leu	Leu	Leu	165	170	175	
Leu	Ser	Ser	Gly	Thr	Ile	Thr	Arg	Val	Ala	Phe	Gly	Lys	Lys	Tyr	Glu	180	185	190	
Gly	Glu	Glu	Glu	Arg	Lys	Asn	Lys	Phe	Ala	Asp	Leu	Ala	Thr	Glu	Leu	195	200	205	
Thr	Thr	Leu	Met	Gly	Ala	Phe	Phe	Val	Gly	Asp	Tyr	Phe	Pro	Ser	Phe	210	215	220	
Ala	Trp	Val	Asp	Val	Leu	Thr	Gly	Met	Asp	Ala	Arg	Leu	Lys	Arg	Asn	225	230	235	240
His	Gly	Glu	Leu	Asp	Ala	Phe	Val	Asp	His	Val	Ile	Asp	Asp	His	Leu	245	250	255	
Leu	Ser	Arg	Lys	Ala	Asn	Gly	Ser	Asp	Gly	Val	Glu	Gln	Lys	Asp	Leu	260	265	270	
Val	Asp	Val	Leu	Leu	His	Leu	Gln	Lys	Asp	Ser	Ser	Leu	Gly	Val	His	275	280	285	
Leu	Asn	Arg	Asn	Asn	Leu	Lys	Ala	Val	Ile	Leu	Asp	Met	Phe	Ser	Gly	290	295	300	
Gly	Thr	Asp	Thr	Thr	Ala	Val	Thr	Leu	Glu	Trp	Ala	Met	Ala	Glu	Leu	305	310	315	320
Ile	Lys	His	Pro	Asp	Val	Met	Glu	Lys	Ala	Gln	Gln	Glu	Val	Arg	Arg	325	330	335	
Val	Val	Gly	Lys	Lys	Ala	Lys	Val	Glu	Glu	Glu	Asp	Leu	His	Gln	Leu	340	345	350	
His	Tyr	Leu	Lys	Leu	Ile	Ile	Lys	Glu	Thr	Leu	Arg	Leu	His	Pro	Val	355	360	365	
Ala	Pro	Leu	Leu	Val	Pro	Arg	Glu	Ser	Thr	Arg	Asp	Val	Val	Ile	Arg	370	375	380	
Gly	Tyr	His	Ile	Pro	Ala	Lys	Thr	Arg	Val	Phe	Ile	Asn	Ala	Trp	Ala	385	390	395	400
Ile	Gly	Arg	Asp	Pro	Lys	Ser	Trp	Glu	Asn	Ala	Glu	Glu	Phe	Leu	Pro	405	410	415	
Glu	Arg	Phe	Val	Asn	Asn	Ser	Val	Asp	Phe	Lys	Gly	Gln	Asp	Phe	Gln	420	425	430	
Leu	Ile	Pro	Phe	Gly	Ala	Gly	Arg	Arg	Gly	Cys	Pro	Gly	Ile	Ala	Phe	435	440	445	
Gly	Ile	Ser	Ser	Val	Glu	Ile	Ser	Leu	Ala	Asn	Leu	Leu	Tyr	Trp	Phe	450	455	460	
Asn	Trp	Glu	Leu	Pro	Gly	Ile										465	470		

-continued

```

<211> LENGTH: 509
<212> TYPE: PRT
<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 43

Met Ala Phe Phe Ser Met Ile Ser Ile Leu Leu Gly Phe Val Ile Ser
1           5           10           15

Ser Phe Ile Phe Ile Phe Phe Phe Lys Lys Leu Leu Ser Phe Ser Arg
           20           25           30

Lys Asn Met Ser Glu Val Ser Thr Leu Pro Ser Val Pro Val Val Pro
           35           40           45

Gly Phe Pro Val Ile Gly Asn Leu Leu Gln Leu Lys Glu Lys Lys Pro
           50           55           60

His Lys Thr Phe Thr Arg Trp Ser Glu Ile Tyr Gly Pro Ile Tyr Ser
65           70           75           80

Ile Lys Met Gly Ser Ser Ser Leu Ile Val Leu Asn Ser Thr Glu Thr
           85           90           95

Ala Lys Glu Ala Met Val Thr Arg Phe Ser Ser Ile Ser Thr Arg Lys
           100          105          110

Leu Ser Asn Ala Leu Thr Val Leu Thr Cys Asp Lys Ser Met Val Ala
           115          120          125

Thr Ser Asp Tyr Asp Asp Phe His Lys Leu Val Lys Arg Cys Leu Leu
           130          135          140

Asn Gly Leu Leu Gly Ala Asn Ala Gln Lys Arg Lys Arg His Tyr Arg
145          150          155          160

Asp Ala Leu Ile Glu Asn Val Ser Ser Lys Leu His Ala His Ala Arg
           165          170          175

Asp His Pro Gln Glu Pro Val Asn Phe Arg Ala Ile Phe Glu His Glu
           180          185          190

Leu Phe Gly Val Ala Leu Lys Gln Ala Phe Gly Lys Asp Val Glu Ser
           195          200          205

Ile Tyr Val Lys Glu Leu Gly Val Thr Leu Ser Lys Asp Glu Ile Phe
210          215          220

Lys Val Leu Val His Asp Met Met Glu Gly Ala Ile Asp Val Asp Trp
225          230          235          240

Arg Asp Phe Phe Pro Tyr Leu Lys Trp Ile Pro Asn Lys Ser Phe Glu
           245          250          255

Ala Arg Ile Gln Gln Lys His Lys Arg Arg Leu Ala Val Met Asn Ala
           260          265          270

Leu Ile Gln Asp Arg Leu Lys Gln Asn Gly Ser Glu Ser Asp Asp Asp
           275          280          285

Cys Tyr Leu Asn Phe Leu Met Ser Glu Ala Lys Thr Leu Thr Lys Glu
290          295          300

Gln Ile Ala Ile Leu Val Trp Glu Thr Ile Ile Glu Thr Ala Asp Thr
305          310          315          320

Thr Leu Val Thr Thr Glu Trp Ala Ile Tyr Glu Leu Ala Lys His Pro
           325          330          335

Ser Val Gln Asp Arg Leu Cys Lys Glu Ile Gln Asn Val Cys Gly Gly
           340          345          350

Glu Lys Phe Lys Glu Glu Gln Leu Ser Gln Val Pro Tyr Leu Asn Gly
           355          360          365

Val Phe His Glu Thr Leu Arg Lys Tyr Ser Pro Ala Pro Leu Val Pro
           370          375          380

Ile Arg Tyr Ala His Glu Asp Thr Gln Ile Gly Gly Tyr His Val Pro
385          390          395          400

```

-continued

Ala Gly Ser Glu Ile Ala Ile Asn Ile Tyr Gly Cys Asn Met Asp Lys
405 410 415

Lys Arg Trp Glu Arg Pro Glu Asp Trp Trp Pro Glu Arg Phe Leu Asp
420 425 430

Asp Gly Lys Tyr Glu Thr Ser Asp Leu His Lys Thr Met Ala Phe Gly
435 440 445

Ala Gly Lys Arg Val Cys Ala Gly Ala Leu Gln Ala Ser Leu Met Ala
450 455 460

Gly Ile Ala Ile Gly Arg Leu Val Gln Glu Phe Glu Trp Lys Leu Arg
465 470 475 480

Asp Gly Glu Glu Glu Asn Val Asp Thr Tyr Gly Leu Thr Ser Gln Lys
485 490 495

Leu Tyr Pro Leu Met Ala Ile Ile Asn Pro Arg Arg Ser
500 505

The invention claimed is:

1. A method for producing a host cell comprising an isoprenoid compound, comprising:

a) providing a host cell that comprises heterologous nucleic acid encoding an isoprenoid synthase, and heterologous nucleic acid encoding one or more protein(s) comprising a cytochrome P450 polypeptide, wherein: the isoprenoid synthase catalyzes production of an isoprenoid compound that is a diterpene or sesquiterpene; and

a cytochrome P450 polypeptide(s) catalyzes hydroxylation, oxidation, demethylation or methylation of a diterpene or sesquiterpene;

b) culturing the host cell under conditions suitable for expressing the isoprenoid synthase and the cytochrome P450 polypeptide(s) and under conditions for producing a diterpene or a sesquiterpene isoprenoid compound not normally produced by the host cell to produce a diterpene or sesquiterpene isoprenoid compound not normally produced by the host cell, wherein the synthase and the cytochrome P450 polypeptide(s) together catalyze the formation of a diterpene or sesquiterpene isoprenoid compound in the host cell, wherein the diterpene or sesquiterpene isoprenoid compound is not normally produced by the cell.

2. The method of claim 1, wherein at least one of the cytochrome P450 polypeptide(s) is a CYP71, CYP73, CYP82 or CYP92 family cytochrome P450.

3. The method of claim 2, wherein each of the CYP71, CYP73, CYP82 and CYP92 family cytochrome P450 polypeptides is encoded by nucleic acid that can be amplified with degenerate primers based on one of SEQ ID NOS: 26-29.

4. The method of claim 1, wherein at least one cytochrome P450 polypeptide has dual hydroxylase activity.

5. The method of claim 1, wherein at least one cytochrome P450 polypeptide has oxidase activity.

6. The method of claim 1, wherein at least one cytochrome P450 polypeptide has hydroxylase activity.

7. The method of claim 1, wherein:
the isoprenoid compound produced by the isoprenoid synthase is a sesquiterpene; and
the isoprenoid synthase is a sesquiterpene synthase.

8. The method of claim 1, wherein:
an isoprenoid compound produced by the isoprenoid synthase is a diterpene; and
the isoprenoid synthase is a diterpene synthase.

9. The method of claim 1, wherein the host cell is a yeast cell, a bacterial cell, an insect cell or a plant cell.

10. The method of claim 1, wherein the host cell is a yeast cell.

11. The method of claim 10, wherein the yeast is *Saccharomyces cerevisiae*.

12. The method of claim 1, wherein at least one of the cytochrome P450 polypeptides is selected from among 5-epi-aristolochene hydroxylase and kaurene oxidase.

13. A host cell, comprising nucleic acid encoding a diterpene or sesquiterpene isoprenoid synthase, and nucleic acid encoding one or more cytochrome P450 polypeptide(s), wherein:

the nucleic acid encoding the synthase and the nucleic acid encoding the P450 polypeptide(s) are heterologous to the host cell;

the synthase catalyzes production of a diterpene or sesquiterpene compound;

the P450 polypeptide(s) catalyze(s) hydroxylation, oxidation, demethylation or methylation of a diterpene or sesquiterpene;

at least one of the P450 polypeptide(s) catalyzes the hydroxylation, oxidation, demethylation or methylation of a diterpene or sesquiterpene whose production is catalyzed by the synthase; and

the host cell thereby can produce a diterpene or sesquiterpene compound not normally produced by the host cell nor by the synthase in the absence of the P450 polypeptide(s).

14. The host cell of claim 13, wherein at least one of the P450 polypeptide(s) catalyzes the dual hydroxylation of a diterpene or sesquiterpene whose production is catalyzed by the synthase.

15. The host cell of claim 13, wherein at least one of the P450 polypeptide(s) is a CYP71, CYP73, CYP82 or CYP92 family cytochrome P450 polypeptide encoded by nucleic acid that can be amplified with degenerate primers based on one of SEQ ID NOS: 26-29.

16. The host cell of claim 13, wherein at least one of the P450 polypeptide(s) comprises a CYP71, CYP73, CYP82 or CYP92 family cytochrome P450 polypeptide.

17. The host cell of claim 13 that is a yeast cell, a bacterial cell, an insect cell or a plant cell.

75

18. The host cell of claim 13 that is a yeast cell.

19. The host cell of claim 13, wherein at least one of the cytochrome P450 polypeptide(s) is selected from among 5-epi-aristolochene hydroxylase and kaurene oxidase.

20. A method for producing an isoprenoid compound, comprising:

a) providing a host cell that comprises heterologous nucleic acid encoding an isoprenoid synthase, and heterologous nucleic acid encoding one or more protein(s) comprising a cytochrome P450 polypeptide, wherein: the isoprenoid synthase catalyzes production of an isoprenoid compound that is a diterpene or sesquiterpene; and

a cytochrome P450 polypeptide(s) catalyzes hydroxylation, oxidation, demethylation or methylation of a diterpene or sesquiterpene;

b) culturing the host cell under conditions suitable for expressing the isoprenoid synthase and the cytochrome P450 polypeptide(s) and under conditions for producing a diterpene or a sesquiterpene isoprenoid compound not normally produced by the host cell to produce a diterpene or sesquiterpene isoprenoid compound not normally produced by the host cell, wherein the synthase and the cytochrome P450 polypeptide(s) together catalyze the formation of the diterpene or sesquiterpene isoprenoid compound; and

c) isolating a diterpene or sesquiterpene isoprenoid compound not normally produced by the cell.

21. A method for producing a host cell comprising an isoprenoid compound, comprising:

a) providing a host cell that comprises heterologous nucleic acid encoding an isoprenoid synthase, and heterologous nucleic acid encoding one or more protein(s) comprising a cytochrome P450 polypeptide, wherein: the isoprenoid synthase catalyzes production of an isoprenoid compound that is a diterpene or sesquiterpene; and

a cytochrome P450 polypeptide(s) catalyzes hydroxylation, oxidation, demethylation or methylation of a diterpene or sesquiterpene;

b) culturing the host cell under conditions suitable for expressing the isoprenoid synthase and the cytochrome P450 polypeptide(s) and under conditions for producing a diterpene or a sesquiterpene isoprenoid compound not normally produced by the host cell to produce a diterpene or sesquiterpene isoprenoid compound not normally produced by the host cell, wherein the synthase and the cytochrome P450 polypeptide(s) together cata-

76

lyze the formation of a diterpene or sesquiterpene isoprenoid compound not normally produced by the cell; and

c) isolating the host cell that comprises an isoprenoid compound not normally produced by the cell.

22. The method of claim 21, further comprising isolating an isoprenoid compound produced by the host cell.

23. The method of claim 1, further comprising isolating an isoprenoid compound that is produced by the host cell.

24. The method of claim 1, further comprising collecting fermentation medium in which the host cell is cultured; and isolating an isoprenoid compound.

25. The method of claim 20, wherein at least one of the cytochrome P450 polypeptide(s) is selected from among 5-epi-aristolochene hydroxylase and kaurene oxidase.

26. A method for producing an isoprenoid compound, comprising:

a) providing a host cell that expresses heterologous nucleic acid encoding a first recombinant protein comprising an isoprenoid synthase, and a second recombinant protein comprising a cytochrome P450 polypeptide, wherein:

the isoprenoid synthase catalyzes production of an isoprenoid compound that is a diterpene or sesquiterpene; and

the cytochrome P450 polypeptide catalyzes hydroxylation, oxidation, demethylation or methylation of the diterpene or sesquiterpene produced by the synthase to produce a diterpene or a sesquiterpene compound not normally produced by the host cell nor whose production is catalyzed by the synthase;

b) culturing the host cell under conditions suitable for expressing the first recombinant protein comprising an isoprenoid synthase and the second recombinant protein comprising the cytochrome P450 polypeptide and under conditions for producing the diterpene or a sesquiterpene isoprenoid compound not normally produced by the host cell, wherein the first and second recombinant protein together catalyze the formation of a diterpene or sesquiterpene isoprenoid compound; and

c) isolating a diterpene or sesquiterpene isoprenoid compound not normally produced by the host cell.

27. The method of claim 26, wherein at least one of the cytochrome P450 polypeptide(s) is selected from among 5-epi-aristolochene hydroxylase and kaurene oxidase.

28. The method of claim 26, wherein the host cell is a yeast cell, a bacterial cell, an insect cell or a plant cell.

29. The method of claim 26, wherein the host cell is a yeast cell.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 8,722,363 B2
APPLICATION NO. : 13/986446
DATED : May 13, 2014
INVENTOR(S) : Chappell et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

IN THE TITLE PAGES:

In Item (71) Applicants, insert:

--Joseph Chappell, Lexington, KY (US)--

Signed and Sealed this
Twenty-ninth Day of July, 2014

A handwritten signature in black ink, reading "Michelle K. Lee". The signature is written in a cursive style with a long, sweeping underline.

Michelle K. Lee
Deputy Director of the United States Patent and Trademark Office